arXiv:1205.1127v1 [math.GR] 5 May 2012

Poincaré Bisectors in Hyperbolic Spaces*

E. Jespers S. O. Juriaans† A. Kiefer A. De A. E Silva
A. C. Souza Filho

Abstract

We determine explicit formulas for the bisectors used in constructing a Dirichlet fundamental domain in hyperbolic two and three space. They are compared with the isometric spheres employed in the construction of a Ford domain and used to find a finite set of generators for discrete groups of finite covolume. Applications are given to Fuchsian groups, Kleinian groups, including the Bianchi groups, and for the construction of a finite set of generators of the unit group of the integral group ring of a finite nilpotent group. An easy implementable algorithm, DAFC, is also given and used in the search for generators of discrete groups.

1 Introduction

Describing generators and relations of groups acting on hyperbolic spaces was started in the nineteenth century. The big difficulty one encounters is the construction of a fundamental domain. This problem was considered by Ford, Poincaré, Serre, Swan, Thurston and many others. Only in the case of a Ford domain explicit formulas are known. Computer aided methods also exist. For Fuchsian groups we refer to [22, 26], for Bianchi groups we refer to [35] and for cocompact groups we refer to [10].

Another non-trivial problem is that of describing units in an order of a non-commutative non-split division algebra, which is related to the problem of finding free pairs in a division algebra. In general, the best one can do is to show the existence of free groups in such orders. This problem is related to the construction of units in the integral group ring $\mathbb{Z}G$ of a finite group G. It is easy to see that $U(\mathbb{Z}G)$, the unit group of $\mathbb{Z}G$, has a subgroup of finite index of the form $\Gamma_1 \times \cdots \times \Gamma_n$, where for each k, $\Gamma_k \subset \mathrm{GL}(n_k, D_k)$, D_k a division algebra. Making use of the congruence theorems, due to Bass-Minor-Serre, Vasserstein-Bak-Rehmann (see [3, 6, 42]), one can construct generators for each Γ_k for many finite groups G. However, there are some exceptional cases that have defied all methods. Some interesting exceptional cases come from fixed point free groups. In this context, in [10], an algorithm is given to compute the unit group of an order in a non-split classical quaternion algebra $\mathcal{H}(K)$ over an imaginary quadratic field extension K of the rationals. The algorithm is

---

*Mathematics subject Classification Primary [$30F40$, $20M25$]; Secondary [$20H10$, $16S34$]. Keywords and phrases: Hyperbolic space, Isometric sphere, Bisector, Bianchi group, Quaternion algebra, Units.
The first author is supported in part by Onderzoeksraad of Vrije Universiteit Brussel and Fonds voor Wetenschappelijk Onderzoek (Flanders). The second author, the corresponding author, was partially supported by CNPq and FAPESP-Brazil, while visiting the Vrije Universiteit Brussel. The third author is supported by Fonds voor Wetenschappelijk Onderzoek (Flanders)-Belgium. The fourth author is supported by FAPESP and CNPq-Brazil. The last author is supported by FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo), Proc. 2008/57930-1.

†Corresponding author.

then applied to the unit group of $\mathcal{H}(\mathbb{Z}[\frac{1+\sqrt{-7}}{2}])$ providing the first example of a description of a finite set of generators of the unit group of a group ring RG matching one of these exceptional cases. Subsequently, in [24], an explicit method was given to construct units in $\mathcal{H}(\mathbb{Z}[\frac{1+\sqrt{-d}}{2}])$ for all $d \equiv 7 \bmod 8$ and it was shown that a unit group of any order of such algebras is a hyperbolic group. However, it remains a challenge to settle the existing exceptional cases.

In this paper our objectives are two fold. The first one is purely theoretical. Making use of the existing theory, we give explicit descriptions of the bisectors in the Poincaré Theory (see [33]) in $\mathbb{H}^n, n \in \{2,3\}$. First these formulas are given in the ball model, where we show that the bisectors are exactly the isometric spheres. Since working in the ball model is not fit for visualization, we deduce explicit formulas of the bisectors in the upper half space (plane) model as well. Equipped with these tools, we turn to describing generators, up to finite index, of a discrete group Γ acting on a hyperbolic 2 or 3-space and having finite covolume (coarea). We give an algorithm, called the Dirichlet Algorithm of Finite Covolume (DAFC for short), to obtain generators for a subgroup of finite index of Γ. The algorithm DAFC is very easy to implement and reduces in much the computational time.

Our next objective is applications. We first revisit a result of [29] which describes those Fuchsian groups having a fundamental domain which is at the same time a Ford and a Dirichlet Domain, called a DF domain. We give an easy and algebraic criterium which readily can be checked once one has a set of generators. Such a criterium also is proved for Kleinian groups. In particular, we prove that the figure-eight knot group and the Whitehead link complement group do not have a DF domain. Next we extend the result of [10] showing that, using our setting, one can obtain generators without the need of a software. We also revisit results from [12, 14] and [26]. For the Bianchi groups, $\mathrm{PSL}(2,\mathfrak{o}_K)$, $K = \mathbb{Q}(\sqrt{-d})$, d a square free positive integer and $\mathfrak{o}_K$ the ring of integers of K, we describe a Poincaré fundamental domain together with its ideal vertices. This is done in a complete way and independently of [40], relying only on one number theoretical result. We also find generators of a subgroup of finite index for these groups. From our description it easily follows that there are no ideal vertices for $d \in \{1,2,3,7,11,19\}$ and that all examples given in [12, Chapter VII] and the case $d = 19$ have a DF domain. We also describe a Poincaré fundamental domain of the figure-eight knot group whose sides are Poincaré bisectors. Finally we give applications to the construction of generators of a subgroup of finite index in the unit group of a group ring.

The outline of the paper is as follows. In Section 2 we record fundamentals of hyperbolic geometry and group rings. In Section 3, which is the main section of the paper, we prove that, in the ball model of hyperbolic space, the bisectors are the isometric spheres, we give explicit formulas and deduce the correspondent formulas in the upper half space (plane) model. We also give conditions for an isometric sphere, in the upper half space (plane) model, to be a bisector and describe symmetries that most of the orders we work with have in common. In Section 4 we handle the DF domains, complementing the results of [29]. We also revisit the results of [10] and describe a Poincaré fundamental domain for the unit group of orders in several division algebras and for the Bianchi groups. As a consequence we obtain generators for these groups. All calculations can be easily verified by the reader. Still in this section, we handle, the ideal vertices of the Bianchi groups, the figure-eight knot group and the Whitehead Link complement group. In the final Section, we describe units, up to finite index, of the unit group of orders in a Wedderburn component of $\mathbb{Q}G$ of exceptional type, for a finite nilpotent group G. We do not rewrite these elements as group ring elements, this can be done easily by using a known complete set of matrix units of these components.

2 Background

In this section we begin by recalling basic facts on hyperbolic spaces and we fix notation. Standard references are [7, 8, 12, 16, 30, 33]. By $\mathbb{H}^n$ (respectively $\mathbb{B}^n$) we denote the upper half space (plane) (respectively the ball) model of hyperbolic n space.

Let K be an algebraic number field and let $\mathfrak{o}_K$ be an order in K. For a and b nonzero elements of K, we denote by $\mathcal{H}(a,b) = \mathcal{H}(a,b,K)$ the generalized quaternion algebra over K, that is $\mathcal{H}(a,b,K) = K[i,\, j : i^2 = a,\, j^2 = b,\, ji = -ij]$. In the particular case in which $a = b = -1$, we simply denote this algebra as $\mathcal{H}(K)$ and as $\mathcal{H}$ if furthermore $K = \mathbb{R}$. The set $\{1,\, i,\, j,\, k = ij\}$ is an additive K-basis of $\mathcal{H}(a,b,K)$. If $a, b \in \mathfrak{o}_K$ then we denote by $\mathcal{H}(a,\, b, \mathfrak{o}_K) = \mathfrak{o}_K + \mathfrak{o}_K i + \mathfrak{o}_K j + \mathfrak{o}_K k$, a subring of $\mathcal{H}(a,b,K)$. By N we denote the usual norm on $\mathcal{H}(a,b,K)$, that is $N(u_0 + u_1 i + u_2 j + u_3 k) = u_0^2 - a u_1^2 - b u_2^2 + ab u_3^2$. In the special case of $\mathcal{H}$ we simply denote $N(x)$ as $|x|$. Denote by $\mathrm{SL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K))$ the multiplicative group $\{x \in \mathcal{H}(a,b,\mathfrak{o}_K) \mid N(x) = 1\}$.

Given $u = u_0 + u_1 + u_2 j + u_3 k \in \mathcal{H}$, let $u' = k^{-1} u k$, $u^* = u_0 + u_1 i + u_2 j - u_3 k$, $\overline{u} = u_0 - u_1 i - u_2 j - u_3 k$. The mapping $u \mapsto u'$ defines an algebra isomorphism of $\mathcal{H}$ and both $u \mapsto u^*$ and $u \mapsto \overline{u}$ define involutions of $\mathcal{H}$. Clearly, $u \in \mathbb{C} + \mathbb{R}j$ if and only if $u^* = u$. The map $\mathcal{H}(a,b,K) \to \mathrm{M}(2,\mathbb{C})$, given by $u = u_0 + u_1 i + u_2 j + u_3 k \mapsto \gamma_u = \begin{pmatrix} u_0 + u_1\sqrt{a} & u_2\sqrt{b} + u_3\sqrt{ab} \\ u_2\sqrt{b} - u_3\sqrt{ab} & u_0 - u_1\sqrt{a} \end{pmatrix}$, is a monomorphism of algebras (see [12, Chapter X]).

Let $\mathbb{H}^3 = \mathbb{C}\times]0,\infty[$. As is common, we shall often think of $\mathbb{H}^3$ as a subset of $\mathcal{H} = \mathcal{H}(-1,-1,\mathbb{R})$ by identifying $\mathbb{H}^3$ with the subset $\{z + rj \in \mathcal{H} : z \in \mathbb{C}, r \in \mathbb{R}^+\} \subset \mathcal{H}$. The ball model $\mathbb{B}^3$ may be identified in the same way with $\{z + rj \in \mathbb{C} + \mathbb{R}j \mid |z|^2 + r^2 < 1\} \subset \mathcal{H}$. Denote by $\mathrm{Iso}(\mathbb{H}^3)$ (respectively $\mathrm{Iso}(\mathbb{B}^3)$) the group of isometries of $\mathbb{H}^3$ (respectively $\mathbb{B}^3$). The groups of orientation preserving isometries are denoted by $\mathrm{Iso}^+(\mathbb{H}^3)$ and $\mathrm{Iso}^+(\mathbb{B}^3)$ respectively. It is well known that $\mathrm{Iso}^+(\mathbb{H}^3)$ and $\mathrm{Iso}^+(\mathbb{B}^3)$ are isomorphic with $\mathrm{PSL}(2,\, \mathbb{C})$ and $\mathrm{Iso}(\mathbb{H}^3)$ and $\mathrm{Iso}(\mathbb{B}^3)$ are isomorphic with $\mathrm{PSL}(2,\, \mathbb{C}) \times C_2$. The action of $\mathrm{PSL}(2,\, \mathbb{C})$ on $\mathbb{H}^3$ is given by $\begin{pmatrix} a & b \\ c & d \end{pmatrix}(P) = (aP + b)(cP + d)^{-1}$, where $(aP + b)(cP + d)^{-1}$ is calculated in $\mathcal{H}$. Explicitly, if $P = z + rj$ and $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then $\gamma(P) = \frac{(az+b)(\overline{cz}+\overline{d}) + a\overline{c}r^2}{|cz+d|^2 + |c|^2 r^2} + (\frac{r}{|cz+d|^2+|c|^2r^2})j$.

The group $\mathrm{SL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K))$ acts as orientation preserving isometries on $\mathbb{H}$ via the embedding $u \mapsto \gamma_u$. The kernel of this action is $\mathrm{I}(a,b,K) = \mathrm{SL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K)) \cap \mathcal{U}(\mathfrak{o}_K)$. We denote by $\mathrm{PSL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K)) = \mathrm{SL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K))/\mathrm{I}(a,b,K)$. Hence finding a set of generators for $\mathrm{SL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K))$ amounts to finding a set of generators for $\mathrm{PSL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K))$.

Let $\mathrm{SB}_2(\mathcal{H}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M(2,\mathcal{H}) \middle|\ d = a',\ b = c',\ a\overline{a} - c\overline{c} = 1 \right\}$. If $f = \begin{pmatrix} a & c' \\ c & a' \end{pmatrix} \in \mathrm{SB}_2(\mathcal{H})$ then $f^{-1} = \begin{pmatrix} \overline{a} & -\overline{c} \\ -c^* & a^* \end{pmatrix}$. The group $\mathrm{SB}_2(\mathcal{H})/\{1,-1\}$ is isomorphic to $\mathrm{Iso}^+(\mathbb{B}^3)$ and hence $\mathrm{SB}_2(\mathcal{H})$ is isomorphic to $\mathrm{SL}(2,\mathbb{C})$ (see [12, Section 1.2]). Let $g = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & j \\ j & 1 \end{pmatrix} \in M_2(\mathcal{H})$. The map $\Psi : \mathrm{SL}(2,\mathbb{C}) \to SB_2(\mathcal{H})$ given by $\Psi(\gamma) = \overline{g}\gamma g$ is a group isomorphism. Explicitly, for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ we have $\Psi(\begin{pmatrix} a & b \\ c & d \end{pmatrix}) = \frac{1}{2} \begin{pmatrix} a + \overline{d} + (b - \overline{c})j & b + \overline{c} + (a - \overline{d})j \\ c + \overline{b} + (d - \overline{a})j & \overline{a} + d + (c - \overline{b})j \end{pmatrix}$ and hence $\|\Psi(\gamma)\|^2 = \|\gamma\|^2 = |a|^2 + |b|^2 + |c|^2 + |d|^2$. The map $\eta_0 : \mathbb{H}^3 \longrightarrow \mathbb{B}^3$, with $P \mapsto (P - j)(-jP + 1)^{-1}$, is an Ψ-equivariant isometry between the hyperbolic models $\mathbb{H}^3$ and $\mathbb{B}^3$, that is $\eta_0(MP) = \psi(M)\eta_0(P)$, for $P \in \mathbb{H}^3$ and $M \in \mathrm{SL}(2,\mathbb{C})$ (see [12, Proposition, I.2.3]).

The following lemma will be useful in the next section. It is a simple consequence of the existence of an inverse element in $SB_2(\mathcal{H})$.

Lemma 2.1 *Let $a, c \in \mathcal{H}$ be such that $a\overline{a} - c\overline{c} = 1$. Then*

1. $c \cdot \overline{a} = a' \cdot c^*$, $a^* c = c^* a$ *and* $a\overline{c} - c' a^* \in \mathbb{R}k$,
2. $\overline{a} \cdot c' = \overline{c} \cdot a'$, $a + a^* \in \mathbb{C} + \mathbb{R}j$
3. $\overline{a}c + \overline{c}a = 2\langle a|c\rangle$, $|a+c|^2 = |a|^2 + |c|^2 + 2\langle a|c\rangle$, *where $\langle \cdot|\cdot\rangle$ is the scalar product in $\mathbb{R}^4$.*

The hyperbolic distance ρ in $\mathbb{H}^3$ is determined by $\cosh \rho(P,P') = \delta(P,P') = 1 + \frac{d(P,P')^2}{2rr'}$, where d is the Euclidean distance and $P = z + rj$ and $P' = z' + r'j$ are two elements of $\mathbb{H}^3$.

Let Γ be a discrete subgroup of $\mathrm{Iso}^+(\mathbb{B}^3)$. The Poincaré method can be used to give a presentation of Γ (see for example [33]). Let Γ_0 be the stabilizer in Γ of $0 \in \mathbb{B}^3$ and let $\mathcal{F}_0$ a fundamental domain for Γ_0. Denote by $D_\gamma(0) = \{u \in \mathbb{B} \mid \rho(0,u) \leq \rho(u,\gamma(0))\}$. Then $\mathcal{F} = \mathcal{F}_0 \cap (\bigcap_{\gamma \in \Gamma \setminus \Gamma_0} D_\gamma(0))$ is a Poincaré fundamental domain of Γ with center 0. Indeed, since Γ is discrete, there exists $P \in \mathbb{B}^3$ such that Γ_P is trivial. Let γ be an isometry mapping 0 to P. It is now easy to see that we get the mentioned fundamental domain from a Poincaré fundamental domain of $\gamma^{-1}\Gamma\gamma$. If Γ is also geometrically finite then a set of generators for Γ consists of the elements $\gamma \in \Gamma$ so that $\mathcal{F} \cap \gamma(\mathcal{F}) \neq \emptyset$ together with Γ_0. In our case $\Gamma = \langle \Gamma_0, \gamma \mid \gamma(\mathcal{F}) \cap \mathcal{F} \neq \emptyset \rangle$ (see [12, Proposition II.8.4]). As a consequence, one obtains generators for $\mathrm{SL}_1(\mathcal{H}(a,\, b, \mathfrak{o}_K))$, by lifting the previous obtained generators and adding the finite set $\mathrm{I}(a, b, K)$.

If S_1 and S_2 are two intersecting spheres in the extended hyperbolic space, then (S_1, S_2) denotes the cosinus of the angle at which they intersect, the dihedral angle. Explicit formulas can be found in [7]. Elements x and y of hyperbolic space are inverse points with respect to S_1 if $y = \sigma(x)$, where σ is the reflection in S_1. In case $S_1 = \partial\mathbb{B}^3 = S^2$, the boundary of $\mathbb{B}^3$, then the inverse point of x with respect to S_1 is denoted by x^*. This should give no confusion with the same notation used in the quaternion algebra and should be clear from the context.

For $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M(2,\mathbb{C})$, we write $a = a(\gamma), b = b(\gamma), c = c(\gamma)$ and $d = d(\gamma)$ when it is necessary to stress the dependence of the entries on the matrix γ. We also have that $\Psi(\gamma) = A_\gamma \sigma_\gamma$, where A_γ is an orthogonal map and σ_γ is the reflection in the isometric sphere of $\Psi(\gamma)$.

We denote by $B_r(p)$, the Euclidean ball with center p and radius r. Its topological closure will be denoted by $\overline{B}_r(p)$. This notation will be used freely during the text.

Let A be a finite dimensional semisimple algebra over the rationals. A subring Γ of A is said to be a ($\mathbb{Z}$)-order if Γ is a finitely generated $\mathbb{Z}$-module that contains a $\mathbb{Q}$-basis of A. It is well known that the unit group $\mathcal{U}(\Gamma)$ is finitely presented. We are mainly interested in the case when $A = RG$, the group ring of a finite group G over the ring of integers R of an algebraic number field (see for example [27, 38]). Finding finitely many generators for $\mathcal{U}(\Gamma)$ is not trivial and is one of the important problems in the field. Nevertheless, for many finite groups G a specific finite set B of generators of a subgroup of finite index in $\mathcal{U}(\mathbb{Z}G)$ has been given.

If $e_1, \ldots, e_n$ denote the primitive central idempotents of the semisimple algebra A then $Ae_i = M(n_i, D_i)$, with D_i a skew field and $n_i \in \mathbb{N}$, are the Wedderburn components of A. If $\emptyset_i$ is an order in D_i then $\prod_i M(n_i, \emptyset_i)$ is an order in A. Hence if $U(\Gamma)$ is the group of units of an order $\Gamma < A$, then $U(\Gamma)$ contains a subgroup of finite index of the form $V_1 \times \cdots \times V_n$, with each V_i a subgroup of finite index in $\mathrm{GL}(n_i, \emptyset_i) = \mathcal{U}(M(n_i, \emptyset_i))$.

Let $\mathrm{SL}(n_i, \emptyset_i)$ denote the group of matrices of reduced norm 1. For an ideal $\mathcal{Q}$ of $\emptyset_i$, let $E(\mathcal{Q}) = \langle I + qE_{lm}, q \in \mathcal{Q}, l \neq m \rangle \leq \mathrm{SL}(n_i, \emptyset_i)$. The celebrated theorems of Bass-Milnor-Serre [6], Vaserstein [42] and Bak-Rehmann [3] state that if $n_i \geq 3$, then $(\mathrm{SL}(n_i, \mathcal{O}_i) : E(\mathcal{Q})) < \infty$ and if $n_i = 2$ and if $\mathcal{O}_i$ is an order in the ring of integers of an algebraic number field which is not rational nor imaginary quadratic extension of $\mathbb{Q}$, then $[\mathrm{SL}(2, \mathcal{O}_i) : E(\mathcal{Q})] < \infty$. For a survey, see [34] and [38]. Moreover, D_i is a totally definite quaternion algebra if and only if $D_i \neq K_i$ and $U(\mathcal{O}_i)$ is finite (see [38]). This shows that to get generators, up to finite index, for the unit group of $\mathbb{Z}G$ of an arbitrary finite group G, one should focus on groups having a representation χ of degree 2 such that $\mathbb{Q}(\chi) \subseteq \mathbb{Q}(\sqrt{-d}), d > 0$ or such that $\mathbb{Q}G$ has a Wedderburn component of the form $M(n, D)$,with D a non-commutative division algebra and $n \in \{1, 2\}$. Among such groups are the fixed point free groups which are exactly the Frobenius complements (see [32, Theorem 18.1.iv]). In particular, the non-abelian nilpotent finite fixed point free groups are $Q_8 \times C_n$, with Q_8 the quaternion group of order 8, C_n a cyclic group of order n and such that 2 has odd multiplicative order in $\mathcal{U}(\mathbb{Z}_n)$.

A Bass cyclic unit is a unit of the form $(1+g+\cdots+g^{i-1})^m + \left(\frac{1-i^m}{o(g)}\right)\widehat{g}$, where $g \in G$, $1 < i < o(g)$, $(i, o(g)) = 1$ and m is the order of i in $\mathcal{U}(\mathbb{Z}_{o(g)})$ (or one takes a fixed multiple of m, for example $\varphi(|G|)$), where φ is the Euler φ-function and $\hat{g} = 1 + g + \cdots + g^{o(g)-1}$. The unipotent units of the form $1 + (1-g)h\widehat{g}$ and $1 + \widehat{g}h(1-g)$, with $g, h \in G$, are called the bicyclic units of $\mathbb{Z}G$.

3 Bisectors and Isometric Spheres

The main purpose of this section is to give explicit formulas for the bisectors in the Poincaré method in two and three dimensional hyperbolic space. To do so, we first prove that, in the ball model, these bisectors are precisely the isometric spheres. Once this has been done, we use simple hyperbolic geometry to give the formulas for the bisectors in the upper half space (plane) model. Calculations are done in dimension three and at the end of the section we also give formulas in dimension two. Standard facts about the theory of hyperbolic geometry will be used freely. Standard references are [7, 8, 12, 16, 33].

Let $0 \in \mathbb{B} = \mathbb{B}^3$ be the origin and $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{C})$ and $\Psi(\gamma) = \begin{pmatrix} A & C' \\ C & A' \end{pmatrix}$.

Recall [12, Section 1.2] that the action of $\Psi(\gamma)$ on $\mathbb{B}^3$ is given by $\Psi(\gamma)(P) = (AP + C')(CP + A')^{-1}$, where the latter is calculated in the classical quaternion algebra $\mathcal{H}(-1, -1, \mathbb{R})$. Consider the Euclidean sphere $\Sigma = \Sigma_{\Psi(\gamma)} = \{P = z + rj \in \mathbb{C} + \mathbb{R}j \mid \|CP + A'\| = 1\}$. It has center $P_{\Psi(\gamma)} = -C^{-1}A'$ and radius $R_{\Psi(\gamma)} = \frac{1}{|C|}$. We remind the reader that * is used both for inverse point with respect to S^2 as well as for the classical involution on the quaternion algebra $\mathcal{H}$.

Also recall that $\gamma \in \mathrm{SU}(2, \mathbb{C})$ if and only if $\gamma(j) = j$ or, equivalently, $\Psi(\gamma)(0) = 0$ (see [7, 12]). In the latter case, we have that $\Psi(\gamma)$ is a linear orthogonal map.

Theorem 3.1 *Let $\gamma \in \mathrm{SL}(2, \mathbb{C})$ with $\gamma \notin \mathrm{SU}(2, \mathbb{C})$. Then $\Sigma_{\Psi(\gamma)} = \{u \in \mathbb{B} \mid \rho(0, u) = \rho(u, \Psi(\gamma^{-1})(0))\}$, the bisector of the geodesic segment linking 0 and $\Psi(\gamma^{-1})(0)$. Moreover $1 + \frac{1}{|C|^2} = |P_{\Psi(\gamma)}|^2$, $D_\gamma(0) = \mathbb{B} \cap Exterior(\Sigma_{\Psi(\gamma)})$ and $P^*_{\Psi(\gamma)} = \Psi(\gamma^{-1})(0)$, inverse points.*

Proof. First, using Lemma 2.1, we have, in $\mathcal{H}$, that $P^*_{\Psi(\gamma)} = P_{\Psi(\gamma)}$, i.e., $P_{\Psi(\gamma)} \in \mathbb{C} + \mathbb{R}j$. As $P_{\Psi(\gamma)} \cdot (\Psi(\gamma^{-1})(0))^{-1} = |A|^2|C|^2 \in \mathbb{R}$, it follows that 0, $P_{\Psi(\gamma)}$ and $\Psi(\gamma^{-1})(0)$ are collinear points. As also $|P_{\Psi(\gamma)}| \cdot |(\Psi(\gamma^{-1})(0))^{-1}| = |-C^{-1}A'| \cdot |-\overline{C}A^{*-1}| = 1$, we have that $P_{\Psi(\gamma)}$ and $\Psi(\gamma^{-1})(0)$ are inverse points with respect to S^2 and hence $(\Sigma, S^2) = 0$, i.e., $1 + R^2_{\Psi(\gamma)} = \frac{1+|C|^2}{|C|^2} = \frac{|A|^2}{|C|^2} = |P_{\Psi(\gamma)}|^2$. It follows also that the reflection, σ say, in Σ is an isometry of $\mathbb{B}$ and that $\sigma \circ \Psi(\gamma^{-1})(0) = 0$. Consequently, $A = \sigma \circ \Psi(\gamma^{-1})$ is an euclidean linear isometry and thus Σ is the isometric sphere of $\Psi(\gamma)$. Here we made use of [7, Theorems 3.4.1, 3.4.2 and 3.5.1].

Now let r be the ray through $P_{\Psi(\gamma)}$ and $M := r \cap \Sigma_{\Psi(\gamma)}$. Clearly $\|M\| = \frac{|A|-1}{|C|}$. Since the hyperbolic metric ρ in $\mathbb{B}$ satisfies $\rho(0,u) = \ln(\frac{1+\|u\|}{1-\|u\|})$ we easily see that $\rho(0,M) = \rho(M, \Psi(\gamma^{-1})(0))$. The ray being orthogonal to Σ, gives us that $\Sigma_{\Psi(\gamma)}$ is the mentioned bisector. ∎

Another way to prove this result is noting that $\phi := \Psi(\gamma)$ is conformal and that $\|d\phi_u \cdot 1\| = \frac{(1-\|\phi(u)\|^2)^2}{(1-\|u\|^2)^2}$ is the scale factor of ϕ_u. Elementary calculations, using Lemma 2.1, show that $\|d\phi_u \cdot 1\| = \frac{1}{|Cu+A'|^2}$. From this we can now read off the equation of the isometric sphere.

Consider $\gamma \in \mathrm{SL}(2,\mathbb{C})$ acting on $\mathbb{H}^3$. We have that $d\gamma_P \cdot 1 = \lim_{t \to 0} \frac{\gamma(P+t)-\gamma(P)}{t}$. Using that $\cosh(d(P,P')) = \delta(P,P') = 1 + \frac{\|P-P'\|^2}{2rr'}$, we obtain that $\|\frac{\gamma(P+t)-\gamma(P)}{t}\| = \frac{1}{\|cP+d\| \cdot \|cP+d+tc\|}$. From this it follows that the scale factor of γ is $\|d\gamma_P \cdot 1\| = \frac{1}{\|cP+d\|^2}$. Hence the isometric sphere of $\gamma \notin \mathrm{SL}(2,\mathbb{C})_\infty$ acting on $\mathbb{H}^3$ is the Euclidean half sphere consisting of the elements $z + rj \in \mathbb{H}^3$ satisfying $|cz+d|^2 + |c|^2 r^2 = 1$. We denote this set by ISO_γ. Its center is denoted by $\hat{P}_\gamma$ and its radius by $\hat{R}_\gamma = \frac{1}{|c|}$.

In general, we do not have that $\mathrm{ISO}_\gamma = \eta_0^{-1}(\Sigma_{\Psi(\gamma)})$, i.e., an isometric sphere in $\mathbb{H}^3$ is not necessarily a Poincaré bisector. For this reason we define $\Sigma_\gamma := \eta_0^{-1}(\Sigma_{\Psi(\gamma)})$. In case it is an Euclidean sphere, we denote its center by P_γ and its radius by R_γ.

Lemma 3.2 *Let* $\gamma \in \mathrm{SL}(2,\mathbb{C})$, *with* $c(\gamma) \neq 0$. *Then*

1. ISO_γ *is the isometric sphere of* $\gamma \in \mathrm{ISO}(\mathbb{H}^3)$.
2. Σ_γ *is the bisector of the geodesic linking* j *and* $\gamma^{-1}(j)$.
3. $j \in \mathrm{ISO}_\gamma$ *if and only if* $|c(\gamma)|^2 + |d(\gamma)|^2 = 1$ *and* $0 \in \mathrm{ISO}_\gamma$ *if and only if* $|d(\gamma)| = 1$.

Proof. (1) was just proved above. (2) follows from Theorem 3.1 and using that η_0 is an isometry between the two models. (3) follows easily. ∎

The following result gives more information on these isometric spheres and bisectors.

Proposition 3.3 *Let* $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{C})$.

1. Σ_γ *is an Euclidean sphere if and only if* $|a|^2 + |c|^2 \neq 1$. *In this case, its center and its radius are respectively given by* $P_\gamma = \frac{-(\overline{a}b+\overline{c}d)}{|a|^2+|c|^2-1}$, $R^2_\gamma = \frac{1+\|P_\gamma\|^2}{|a|^2+|c|^2}$.

2. Σ_γ *is a plane if and only if* $|a|^2 + |c|^2 = 1$. *In this case* $Re(\overline{v}z) + \frac{|v|^2}{2} = 0, z \in \mathbb{C}$ *is a defining equation of* Σ_γ, *where* $v = \overline{a}b + \overline{c}d$.

3. $|\overline{a}b+\overline{c}d|^2 = (|a|^2+|c|^2)(|b|^2+|d|^2)-1$

4. *Suppose* $c \neq 0$*. Then* $|\hat{P}_\gamma - P_\gamma| = \frac{|d-\overline{a}|}{|c|(|a|^2+|c|^2-1)}$*. Moreover* $\mathrm{ISO}_\gamma = \Sigma_\gamma$ *if and only if* $d = \overline{a}$*. In this case we also have that* $c = \lambda\overline{b}$*, with* $\lambda \in \mathbb{R}$*. If* $c = 0$ *and* $\infty \in \Sigma_\gamma$ *then the same conclusion holds.*

5. *Suppose that* $ISO_\gamma = \Sigma_\gamma$ *or that* $c = 0$ *and* $\infty \in \Sigma_\gamma$*. Then* $\mathrm{tr}(\gamma) \in \mathbb{R}$

Proof. Let $P^*_{\Psi(\gamma)} = \Psi(\gamma)^{-1}(0)$. Since η_0 is a Möbius transformation, it follows that $j = \eta_0^{-1}(0)$ and $\gamma^{-1}(j) = \eta_0^{-1}(P^*_{\Psi(\gamma)})$ are inverse points with respect to Σ_γ. So if, Σ_γ is not a vertical plane, then j, $\gamma^{-1}(j)$ and P_γ are collinear points. It follows that $P_\gamma = l \cap \partial\mathbb{H}$, where l is the line determined by j and $\gamma^{-1}(j)$. Since $\gamma^{-1}(j) = -\frac{(\overline{a}b+\overline{c}d)}{|a|^2+|c|^2} + \frac{1}{|a|^2+|c|^2}j$, a simple calculation gives the formula of P_γ. To find the radius of Σ_γ one just has to notice that $R^2_\gamma = \|j - P_\gamma\| \cdot \|\gamma^{-1}(j) - P_\gamma\|$. This proves the first item.

The expression of P_γ shows that Σ_γ is a vertical plane if and only if $|a|^2+|c|^2 = 1$. In this case, $\gamma^{-1}(j) = -(\overline{a}b+\overline{c}d)+j$ and hence $v = j - \gamma^{-1}(j) = \overline{a}b+\overline{c}d$ is orthogonal to Σ_γ. From this one obtains the mentioned defining equation of Σ_γ, hence the second item. The third item is straight forward.

Since $\det(\gamma) = 1$ we have that $|\hat{P}_\gamma - P_\gamma| = |\frac{d}{c} - \frac{\overline{a}b+\overline{c}d}{|a|^2+|c|^2-1}| = \frac{|d-\overline{a}|}{|c|(|a|^2+|c|^2-1)}$. Hence if $\mathrm{ISO}_\gamma = \Sigma_\gamma$ then $d = \overline{a}$ and hence $bc = |a|^2 - 1 \in \mathbb{R}$. With the formula of R_γ at hand, we readily find that $R_\gamma = \frac{1}{|c|}$. In case $c = 0$ and Σ_γ is a plane, we have that $ad = 1$ and $1 = |a|^2+|c|^2 = |a|^2$, and $d = \overline{a}$. If $\det(\gamma) = -1$ we proceed in the same way to obtain that $d = -\overline{a}$. This proves the fourth item.

To prove the last item, use item four. We have that $tr(\gamma) = a + \overline{a} \in \mathbb{R}$. ∎

Proposition 3.4 *Let* $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{C})$ *and* $\Psi(\gamma) = \begin{pmatrix} A & C' \\ C & A' \end{pmatrix}$*. Then the following properties hold.*

1. $\mathrm{ISO}_{\Psi(\gamma)} = \Sigma_{\Psi(\gamma)}$
2. $|A|^2 = \frac{2+\|\gamma\|^2}{4}$, $|C|^2 = \frac{\|\gamma\|^2-2}{4}$ *and* $|A|^2 - |C|^2 = 1$
3. $P_{\Psi(\gamma)} = \frac{1}{-2+\|\gamma\|^2} \cdot [\ -2(\overline{a}b+\overline{c}d) + [(|b|^2+|d|^2) - (|a|^2+|c|^2)]j\ ]$
4. $\Psi(\gamma^{-1})(0) = P^*_{\Psi(\gamma)} = \frac{1}{2+\|\gamma\|^2} \cdot [\ -2(\overline{a}b+\overline{c}d) + [(|b|^2+|d|^2) - (|a|^2+|c|^2)]j\ ]$ *(notation of inverse point w.r.t.* S^2*).*
5. $\|P_{\Psi(\gamma)}\|^2 = \frac{2+\|\gamma\|^2}{-2+\|\gamma\|^2}$
6. $R^2_{\Psi(\gamma)} = \frac{4}{-2+\|\gamma\|^2}$
7. $\Sigma_{\Psi(\gamma)} = \Sigma_{\Psi(\gamma_1)}$ *if and only if* $\gamma_1 = \gamma_0\gamma$*,* $\gamma_0 \in \mathrm{SU}(2,\mathbb{C})$*.*

Proof. The first item was already proved in Theorem 3.1. The proof of the next five items is straightforward using the explicit formulas for $\Psi(\gamma)$, A, C (see Section 2.1) and knowing that $P_{\Psi(\gamma)} = C^{-1}A'$.

To prove the last item. Recall that $\Psi(\gamma_0)(0) = 0$ if and only if $\Psi(\gamma_0)$ is an orthogonal map and that $\Psi(\gamma) \circ \sigma_\gamma$ also is an orthogonal map. From these comments the last item follows readily. ∎

If $\Gamma \subset Iso^+(\mathbb{B}^3)$ is a discrete subgroup of $\mathrm{SL}(2,\mathbb{C})$ then, for each $\lambda \in \mathbb{R}$, the set $\{\gamma \in \Gamma |\ \|\gamma\| = \lambda\}$ is finite (this follows also easily from item 5 of the previous result) and hence, by the Proposition 3.4, for any sequence $(\gamma_n)_n \subset \Gamma$, of two by two distinct elements of Γ, we must have that $R_{\gamma_n} \longrightarrow 0$ as $n \longrightarrow \infty$. Also if $\gamma_1, \gamma \in \Gamma$ have the same norm then, by item 5 of the previous Proposition, the intersection of their isometric sphere with S^2 have the same Euclidean volume. However it is not clear that this volume is a strictly decreasing function of the norm. We shall now address this problem.

Let $\gamma \in \mathrm{SL}(2,\mathbb{C})$ and $\gamma \notin \mathrm{SU}(2,\mathbb{C})$ and let r be the ray through the center of $\Sigma_{\Psi(\gamma)}$. Denote by M and N, respectively, the intersection of r with $\Sigma_{\Psi(\gamma)}$ and S^2. Put $\rho_\gamma = \|\vec{MN}\|$. Explicitly we have that $\rho_\gamma = 1 + R_{\Psi(\gamma)} - \|P_{\Psi(\gamma)}\|$. Our next result shows that ρ_γ is a strictly decreasing function of $\|\gamma\|$. Note that the Euclidean volume of the intersection of $\Sigma_{\Psi(\gamma)}$ with S^2, mentioned above, is a function of ρ_γ.

Lemma 3.5 *Let $\Gamma < \mathrm{PSL}(2,\mathbb{C})$ be a discrete subgroup acting on $\mathbb{B}^3$. Then ρ_γ is a strictly decreasing function of $\|\gamma\|$ on $\Gamma \setminus \Gamma_0$.*

Proof. Let $f(x) = 1 - (\frac{x^2+2}{x^2-2})^{\frac{1}{2}} + 2(x^2-2)^{\frac{-1}{2}}$. Then $f'(x) = -2x(x^2-2)^{-3/2}(x^2+2)^{-1/2}[-2+\sqrt{x^2+2}]$. For any $\gamma \in \mathrm{GL}(2,\mathbb{C})$ we have that $2 \cdot |\det(\gamma)| \le \|\gamma\|^2$. From these facts the result follows. ∎

The following result is very useful if one has to decide if a given group is cocompact or whether a certain point of S^2 is an ideal point. In fact, in Section 4 we use it to show that the Bianchi groups are not cocompact. It is also useful when implementing the DAFC for a given group.

Lemma 3.6 *Let* $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{C})$. *Then*

1. $0 \notin \Sigma_{\Psi(\gamma)}$.
2. $j \in \Sigma_{\Psi(\gamma)}$ *if and only if* $|a|^2 + |c|^2 = 1$.
3. $j \in \mathrm{Interior}(\Sigma_{\Psi(\gamma)})$ *if and only if* $|a|^2 + |c|^2 < 1$.
4. $-j \in \Sigma_{\Psi(\gamma)}$ *if and only if* $|b|^2 + |d|^2 = 1$.
5. $-j \in \mathrm{Interior}(\Sigma_{\Psi(\gamma)})$ *if and only if* $|b|^2 + |d|^2 < 1$.

Proof. All items follow readily if one uses the previous proposition and the explicit expression of $\Psi(\gamma)$. ∎

Making use of $\eta_0 : \mathbb{H}^3 \to \mathbb{B}^3$, the results proved so far in the ball model can be transfered to the upper half space model of hyperbolic space. Note that $\eta_0^{-1}(-j) = 0$, $\eta_0^{-1}(0) = j$, $\eta_0^{-1}(j) = \infty$ and that η_0^{-1} sends the unit disc to $S^2 \cap \mathbb{H}^3$. We will make use of the results of this section in both models, according to which is more convenient.

Corollary 3.7 *Let Γ be a discrete subgroup of* $\mathrm{PSL}(2,\mathbb{C})$ *and* $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. *Suppose that* $\{|a|^2, |b|^2, |c|^2, |d|^2\} \subset \mathbb{N}$. *Then* $\{-j, j\} \cap \mathrm{Interior}(\Sigma_{\Psi(\gamma)}) = \emptyset$. *In particular, if all $\gamma \in \Gamma$ have this property then Γ is not cocompact.*

In the upper half space model, under the assumptions of the corollary and supposing that $\Gamma_0 = 1$, it means that 0 and ∞ are ideal points.

Our next result will be useful when dealing with ideal vertices of Kleinian groups, in particular the Bianchi groups. A version for Fuchsian groups can also be given. We leave this to the reader.

Lemma 3.8 *Let* $\gamma \in \mathrm{PSL}(2,\mathbb{C})$ *and suppose that* $j \in \mathrm{Exterior}(\Sigma_{\Psi(\gamma)}) \cap \mathrm{Exterior}(\Sigma_{\Psi(\gamma^{-1})})$. *Then* $\eta_0(\hat{P}_\gamma) \in \mathrm{Interior}(\Sigma_{\Psi(\gamma)})$.

Proof. We have that $\Psi(\gamma)(\Sigma_{\Psi(\gamma)}) = \Sigma_{\Psi(\gamma^{-1})}$ and the latter disconnects $\mathbb{B}^3$ in two connected components. Since $\Psi(\gamma)(P_{\Psi(\gamma)}) = \infty$ it follows that $\Psi(\gamma)$ maps $\mathrm{Interior}(\Sigma_{\Psi(\gamma)})$ onto $\mathrm{Exterior}(\Sigma_{\Psi(\gamma^{-1})})$. Now observe that $\Psi(\gamma)(\eta_0(\hat{P}_\gamma)) = j$. Hence, because of the hypothesis, the conclusion follows. ∎

Note that in $\mathbb{H}^3$ the hypothesis is that S^2 is not contained in $\mathrm{Interior}(\Sigma) \cap \mathrm{Interior}(\Sigma^{-1})$. In particular, this is true if $R_\gamma < 1$.

The following result is useful if one is interested in the dihedral angles. Making use of the previous results, these can now be calculated knowing only the norm of the two elements involved. Since all models of hyperbolic space of the same dimension are isometric, the formulas of our next result hold for all models.

Lemma 3.9 *Let* $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{C})$, $\pi_0 = \partial\mathbb{H}^n$, $n = 2, 3$, *and θ the angle between $\Sigma_{\Psi(\gamma)}$ and $\Sigma \in \{\Sigma_{\Psi(\gamma_1)}, \pi_0\}$ with $\Sigma \cap \Sigma_{\Psi(\gamma)} \neq \emptyset$. Then*

1. *If* $\Sigma = \Sigma_{\Psi(\gamma_1)}$ *then* $\cos(\theta) = \frac{2|1-\langle P_{\Psi(\gamma)} | P_{\Psi(\gamma_1)}\rangle|}{2R^2_{\Psi(\gamma_1)} \cdot R^2_{\Psi(\gamma)}}$.
2. *If* $\Sigma = \pi_0$ *then* $\cos(\theta) = \frac{|\,|b|^2+|d|^2-|a|^2-|c|^2|}{2\sqrt{\|\gamma\|^2-2}}$

Proof. This follows from the known formulas for these angles and the results of this section (see for example [7, Section 3.2]). ∎

Lemma 3.10 *Let* $\gamma, \gamma_1 \in \mathrm{SL}(2,\mathbb{C})$ *and* $\gamma_2 \in \mathrm{SU}(2,\mathbb{C})$. *Then*

1. $\hat{P}_{\gamma_2\gamma\gamma_1} = \hat{P}_{\gamma\gamma_1} = \gamma_1^{-1}(\hat{P}_\gamma)$
2. $P_{\Psi(\gamma_2\gamma\gamma_1)} = P_{\Psi(\gamma\gamma_1)} = \Psi(\gamma_1^{-1})(P_{\Psi(\gamma)})$

Proof. Note that $P_{\Psi(\gamma)} = \Psi(\gamma^{-1})(\infty)$ and $\hat{P}_\gamma = \gamma^{-1}(\infty)$. Two isometries have identical isometric spheres if and only if they differ by an element of $\mathrm{SU}(2,\mathbb{C})$. These two observations prove the lemma. ∎

We briefly consider the two dimensional model. Let $\gamma \in \mathrm{SL}(2,\mathbb{R})$. As a Möbius transformation, γ acts on $\mathbb{H}^2$ and $\gamma_0 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix}$ gives an isometry between the two models. Proceeding as in the 3-dimensional model, we obtain explicit formulas for the bisectors. We use the same notation as before. Note that the role of j is now played by i and, since we are in the commutative setting, calculations are easier.

Proposition 3.11 *Let* $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{R})$ *and* $\Psi(\gamma) = \gamma_0\gamma\gamma_0^{-1} = \begin{pmatrix} A & \overline{C} \\ C & \overline{A} \end{pmatrix}$ *acting on* $\mathbb{B}^2$. *Then*

1. $\mathrm{ISO}_{\Psi(\gamma)} = \Sigma_{\Psi(\gamma)}$
2. $|A|^2 = \frac{\|\gamma\|^2+2}{4}$, $|C|^2 = \frac{-2+\|\gamma\|^2}{4}$ *and* $|A|^2 - |C|^2 = 1$
3. $P_{\Psi(\gamma)} = \frac{1}{-2+\|\gamma\|^2} \cdot [\ -2(ab+cd) + [(b^2+d^2)-(a^2+c^2)]i\ ]$
4. $\Psi(\gamma^{-1})(0) = P^*_{\Psi(\gamma)} = \frac{1}{\|\gamma\|^2+2} \cdot [\ -2(ab+cd) + [(b^2+d^2)-(a^2+c^2)]i\ ]$ *(notation of inverse point w.r.t.* S^1*).*
5. $\|P_{\Psi(\gamma)}\|^2 = \frac{2+\|\gamma\|^2}{-2+\|\gamma\|^2}$
6. $R^2_{\Psi(\gamma)} = \frac{4}{-2+\|\gamma\|^2}$

In what follows the transformation γ is an element of a Kleinian group. Similar results can be obtained in the Fuchsian case. We establish some results which reveal the relative position of the isometric spheres of γ and γ^{-1}..

Lemma 3.12 *Let* $\gamma \in \mathrm{PSL}(2,\mathbb{C})$ *be a parabolic element and* $z_0 \in \mathbb{C}$ *the fixed point of* γ. *Then* $z_0 = \frac{a-d}{2c} \in \mathbb{Q}(a(\gamma), c(\gamma), d(\gamma))$ *and* $\Sigma_\gamma \cap \Sigma_{\gamma^{-1}} = \{z_0\}$, *i.e.,* Σ_γ *and* $\Sigma_{\gamma^{-1}}$ *are tangent at* z_0.

Proof. Let $a = a(\gamma), b = b(\gamma), c = c(\gamma)$ and $d = d(\gamma)$ and note that $c \neq 0$. Since $\gamma(z_0) = z_0$ we have that $z_0(cz_0+d) = az_0+b$ and hence $cz_0^2 + (d-a)z_0 - b = 0$. The discriminant of this quadratic equation is equal to $(a-d)^2 + 4bc = (a+d)^2 - 4 = tr(\gamma)^2 - 4 = 0$ and hence $z_0 = \frac{a-d}{2c}$.

To prove the second part, let $p = \eta_0(z_0)$ and choose an orientation preserving orthogonal map sending p to j. Such a map is of the form $\Psi(\gamma_0)$, with $\gamma_0 \in \mathrm{SU}(2,\mathbb{C})$. Let $\gamma_1 = \gamma_0\gamma\gamma_0^{-1}$ and note that $\Psi(\gamma_1)(j) = j$. By Proposition 3.4, and the definition of the bisector, we have that $\Sigma_{\Psi(\gamma_0\gamma\gamma_0^{-1})} = \Sigma_{\Psi(\gamma\gamma_0^{-1})} = \Psi(\gamma_0)(\Sigma_{\Psi(\gamma)})$. In $\mathbb{H}^3$ we have that $\gamma_1(\infty) = \infty$ and hence $c(\gamma_1) = 0$ and $a(\gamma_1) = d(\gamma_1) = 1$. Using Proposition 3.3 we obtain that Σ_{γ_1} and $\Sigma_{\gamma_1^{-1}}$ are parallel planes and hence they intersect only at ∞. Since $\Psi(\gamma_0)$ is an orthogonal map it follows that $\Sigma_{\Psi(\gamma)}$ and $\Sigma_{\Psi(\gamma^{-1})}$ are parallel bisectors and thus $\Sigma_\gamma \cap \Sigma_{\gamma^{-1}} = \{z_0\}$. ∎

Lemma 3.13 *Let* $\gamma \in \mathrm{PSL}(2,\mathbb{C})$ *be a hyperbolic or loxodromic element and* $z_0,\ z_1 \in \partial(\mathbb{B}^3)$ *the two fixed points of* γ. *Then the following properties hold.*

1. $\{z_0, z_1\} \subset \mathrm{Interior}(\Sigma_{\Psi(\gamma)}) \cup \mathrm{Interior}(\Sigma_{\Psi(\gamma^{-1})})$.
2. *If* γ *is hyperbolic then* $z_0 \in \Sigma_{\Psi(\gamma)}$, $z_1 \in \Sigma_{\Psi(\gamma^{-1})}$ *and* $\Sigma_{\Psi(\gamma)} \cap \Sigma_{\Psi(\gamma^{-1})} = \emptyset$.
3. *If* γ *is loxodromic then there exists* $n \in \mathbb{N}$ *such that* $\Sigma_{\Psi(\gamma^n)} \cap \Sigma_{\Psi(\gamma^{-n})} = \emptyset$.

Proof. Choose $\gamma_0 \in \mathrm{SU}(2,\mathbb{C})$ such that $\Psi(\gamma_0)(j) = z_0$ and set $\gamma_1 = \gamma_0\gamma\gamma_0^{-1}$. Let $a = a(\gamma_1)$, $b = b(\gamma_1)$, $c = c(\gamma_1)$ and $d = d(\gamma_1)$. Then $c(\gamma_1) = 0$ and $|a(\gamma_1)| \neq 1 \neq |d(\gamma_1)|$. Switching to γ_1^{-1} if necessary, we may suppose that $|a(\gamma_1)| < 1$ and hence, by Lemma 3.6, $j \in \mathrm{Interior}(\Sigma_{\Psi(\gamma_1)})$. Since $\Sigma_{\Psi(\gamma_0\gamma\gamma_0^{-1})} = \Psi(\gamma_0)(\Sigma_{\Psi(\gamma)})$, it follows that $z_0 \in \mathrm{Interior}(\Sigma_{\Psi(\gamma)})$.

Working in $\mathbb{H}^3$, we have that ∞ and $\frac{b}{d-a} = \frac{ab}{1-a^2}$ are the fixed points of γ_1, $P_{\gamma_1} = \frac{\overline{a}b}{1-|a|^2}$, $P_{\gamma_1^{-1}} = \frac{a}{\overline{a}}P_{\gamma_1}$, $R^2_{\gamma_1} = \frac{(1-|a|^2)^2+|ab|^2}{|a|^2(1-|a|^2)^2}$ and $R_{\gamma_1^{-1}} = |a|^2R_{\gamma_1}$. From this it follows that $R_{\gamma_1} - (|P_{\gamma_1} - P_{\gamma_1^{-1}}| + R_{\gamma_1^{-1}})=(1-|a|^2)R_{\gamma_1} - \frac{|\overline{a}-a|}{|a|}|P_{\gamma_1}|$. The condition we look for is that $0 < (1-|a|^2)R_{\gamma_1} - \frac{|\overline{a}-a|}{|a|}|P_{\gamma_1}|$. Using the expressions obtained, it follows that $|P_{\gamma_1}|^2[\frac{4\mathrm{Im}^2(a)-(1-|a|^2)^2}{|a|^2}] < \frac{(1-|a|^2)^2}{|a|^2}$. If γ is hyperbolic then, since $c(\gamma) = 0$ we have that $a \in \mathbb{R}$, the left hand side is negative and so the bisectors are disjoint. Also, still in the hyperbolic case, $P_{\gamma_1} = z_1$. Noting that $a(\gamma^n) = (a(\gamma))^n$, it follows that, in the loxodromic case, we may choose n such that the left had side is negative. ▮

Working in $\mathbb{H}^3$ it is easy to see that the axis of γ is perpendicular to both Σ_γ and $\Sigma_{\gamma^{-1}}$ if and only if z_0 and z_1 are antipodal points in $\mathbb{B}^3$. The following corollary is [33, Corollary 1 of Theorem XII.2.4] in case the element γ is hyperbolic.

Corollary 3.14 *Let* Γ *be a discrete group acting on* $\mathbb{B}^3$*,* $\mathcal{F}$ *a fundamental domain of* Γ*,* $\gamma \in \Gamma$ *and* z_0 *a fixed point of* γ*. If* γ *is hyperbolic or loxodromic then* $z_0 \notin \partial\mathcal{F}$*.*

Let γ be an elliptic element such that $c = c(\gamma) = 0$. The fixed points of γ in $\mathbb{H}^3$ are ∞ and $z_1 = \frac{bi}{2\mathrm{Im}(a)}$. It follows by Proposition 3.3 that Σ_γ and $\Sigma_{\gamma^{-1}}$ are vertical planes that intersect in the vertical line l, say, through z_1. Making use of Lemma 2.1 and Proposition 3.3, we find that the angle of rotation of γ around l is $\frac{\mathrm{Re}(a^2)}{|a|^2}$. Hence we obtain the following.

Lemma 3.15 *Let* γ *be an elliptic element of* $\mathrm{PSL}(2,\mathbb{C})$*. Then* $\Sigma_{\Psi(\gamma)} \cap \Sigma_{\Psi(\gamma^{-1})}$ *is a circle, containing both fixed point of* γ*.*

In the next section we shall give generating sets of some discrete groups Γ. Since some of these sets are large, in order to write these in a more compact form, we look for symmetries of the fundamental domain of Γ. We proceed to describe some general isomorphism and involutions of $\mathrm{PSL}(2,\mathbb{C})$.

Let $\gamma \in \mathrm{SL}(2,\mathbb{C})$. Denote by σ conjugation by the matrix $\begin{pmatrix} \sqrt{i} & 0 \\ 0 & \sqrt{-i} \end{pmatrix}$, by δ we denote conjugation by the matrix $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\tau(\gamma) = \overline{\gamma}$, denotes complex conjugation of the entries of γ. If no confusion arises, σ will also denote the matrix $\begin{pmatrix} \sqrt{i} & 0 \\ 0 & \sqrt{-i} \end{pmatrix}$. It is easily seen that $P_{\sigma(\gamma)} = iP_\gamma$, $R_{\sigma(\gamma)} = R_\gamma$, $P_{\tau(\gamma)} = \overline{P_\gamma}$ and $R_{\tau(\gamma)} = R_\gamma$. Let $\phi = \sigma^2 \circ \delta \circ \tau$. Then, working in the ball model, it follows easily that $P_{\Psi(\phi(\gamma))} = \pi(P_{\Psi(\gamma)})$, where π is the reflection in the xy-plane. Since $\eta_0 : \mathbb{H}^3 \to \mathbb{B}^3$ is equivariant, it follows that $P_{\phi(\gamma)}$ is the reflection of P_γ in S^2, but the radius is not maintained. Note that $\sigma^2 \circ \tau$, preserves the radius of bisectors and, as a symmetry, corresponds to the reflection in the imaginary axis of the complex plane: $P_{\sigma^2\circ\tau(\gamma)} = -\overline{P_\gamma}$. The maps τ and σ^2 induce isometries of $\mathbb{H}^n$, $n \in \{2,3\}$. The first is reflexion in the real axis in case $n = 2$ or in the plane spanned by 1 and j in case $n = 3$ and the second is the reflection in the origin. The isometries σ^2 and $\sigma^2 \circ \tau$ satisfy $\sigma^2(z + rj) = -z + rj$, $\sigma^2 \circ \tau(z + rj) = -\overline{z} + rj$, $z \in \mathbb{C}$ and $r > 0$.

In Section 4.2, all our examples are division quaternion algebras, $\mathcal{H}(a, b, \mathbb{Q}(\sqrt{-d}))$, with $d > 0, a, b \in \mathbb{R}, ab > 0$. The orders, in the algebras we work with, induce tessellations on $\mathbb{H}^3$ which are invariant under some subgroup generated by the symmetries defined above. These symmetries, induce algebra isomorphisms on $\mathcal{H}(a, b, \mathbb{Q}(\sqrt{-d}))$, all of which can be written explicitly. We shall use the same notation for a symmetry on $\mathbb{H}^3$ as for the induced algebra isomorphism.

Let $\Gamma < \mathrm{SL}(2, \mathbb{C})$ be a discrete group, $\gamma \in \Gamma, 1 \neq \gamma_1 \in \Gamma_\infty$ and Σ_{γ_1} a vertical plane. Since $d(\gamma) = \overline{a(\gamma)}$ has norm 1, it follows that $P_{\gamma\gamma_1} = \gamma_1^{-1}(P_\gamma)$ and that γ_1 acts as an Euclidean isometry on $\partial\mathbb{H}^3$ and each horosphere of $\mathbb{H}^3$. Hence $\gamma_1^{-1}(\Sigma_\gamma)$ is the sphere with center $P_{\gamma\gamma_1}$ and whose radius is equal to R_γ. This shows that if a point $z \in \mathbb{C}$ is not covered by the tessellation induced by Γ, then $\gamma_1(z)$ is also not covered by this tessellation and hence, in case Γ is of finite covolume, it has a finite number of ideal vertices $\{z_1, \cdots z_n\}$ such that a point $z \in \mathbb{C}$ is not covered by the tessellation induced by Γ if and only if $z \in \bigcup_{1 \leq k \leq n} \Gamma_\infty \cdot z_k$.

In the case of a Ford domain, the same proof of the previous paragraph shows that the set of singular points is invariant under the action of Γ_∞ (see for example [12, Proposition VII.2.9]).

Let $\gamma \in \mathrm{PSL}(2, \mathbb{C})$. It is known that $\Psi(\gamma)(\Sigma_{\Psi(\gamma)}) = \Sigma_{\Psi(\gamma^{-1})}$ and hence, using the definition of Σ_γ and that η_0 is $\mathrm{SL}(2, \mathbb{C})$-equivariant, we have that $\gamma(\Sigma_\gamma) = \Sigma_{\gamma^{-1}}$. It is also known that $\gamma(ISO_\gamma) = ISO_{\gamma^{-1}}$, if $c(\gamma) \neq 0$. Of course, this is now surprise since Σ_γ is a Poincaré biseсctor.

Another useful fact is that $P_{\Psi(\gamma)} = P_{\Psi(\gamma^{-1})}$ if and only if $\gamma^2 = 1$ and this is true if and only if $d(\gamma) = -a(\gamma)$. The latter happens if and only if $\langle A_\gamma, \sigma_\gamma \rangle$ is a dihedral group. The reader should compare some results and expressions proved here with some results and expressions of [7, 33], especially for $\mathbb{H}^2$.

We now have covered enough of a Ford Theory for Kleinian groups. From the results of this section, it follows that this is the same as the Poincaré Theory. In the next section we shall apply this theory, thus showing its usefulness and consistency.

4 Applications

In this section all groups $\Gamma < \mathrm{PSL}(2, \mathbb{C})$ are discrete groups and are either Fuchsian or Kleinian. If Γ is cocompact, then it is well known that it contains no parabolic elements. We provide a proof of this result, making use of Lemma 3.9. Indeed, if $\gamma_0 \in \Gamma$ is parabolic then, working with a conjugate of Γ if necessary, we may suppose that $\gamma_0(z) = z + 1$. Hence, by Shimizu´s Lemma (see [12, Theorem II.3.1]), for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma, \gamma \notin \Gamma_\infty$, we have that $|c| \geq 1$. It follows that $|a|^2 + |c|^2 \geq 1$ and thus, $j \notin \mathrm{Interior}(\Sigma_{\Psi(\gamma)})$. If $\gamma \in \Gamma_\infty$ then, by [12, Theorem II.1.8], $|a| = 1$ and hence $j \in \Sigma_{\Psi(\gamma)}$. It follows that j is an ideal vertex of Γ, a contradiction.

Lemma 4.1 *Let $\Gamma < \mathrm{PSL}(2, \mathbb{C})$ act on $\mathbb{H}^3$. Then the following are equivalent.*

1. *Γ is discontinuous.*
2. *If (γ_n) is a sequence of distinct elements in Γ, then $(\Psi(\gamma_n^{-1})(0))$ has no limit point in $\mathbb{B}^3$.*
3. *If (γ_n) is a sequence of distinct elements in Γ, then $\lim_{n\to\infty} \|\gamma_n\| = \infty$.*
4. *Γ is discrete.*

Proof. We first prove that (1) and (2) are equivalent. Suppose that Γ is discontinuous and let $(\gamma_n) \subset \Gamma$ be a sequence of distinct elements. Then, by hypothesis, $(\Psi(\gamma_n^{-1})(0))$ has no limit point in $\mathbb{B}^3$. Suppose now that if $(\gamma_n) \subset \Gamma$ is a sequence of distinct elements, $(\Psi(\gamma_n^{-1})(0))$ has no limit point in $\mathbb{B}^3$. By Proposition 3.4, it follows that $\|\Psi(\gamma_n^{-1})(0)\| \to 1$ and hence $\lim\limits_{n\to\infty} \|\gamma_n\| = \infty$. Choose $P \in \mathbb{B}^3$ and let $n_0 \in \mathbb{N}$ be such that $P \in \text{Exterior}(\Sigma_{\Psi(\gamma_n)}), n \geq n_0$. Then $\gamma_n(P) \in \text{Interior}(\Sigma_{\Psi(\gamma_n^{-1})})$ and since $R_{\Psi(\gamma_n)} \to 0$, it follows that $(\gamma_n(P))$ has no limit point in $\mathbb{B}^3$. Note that we made use repeatedly of Proposition 3.4.

The fact that item (2) and (3) are equivalent follows from Proposition 3.4.
Suppose that Γ is discrete and let $(\gamma_n) \subset \Gamma$ be a sequence of distinct elements. Suppose by contradiction, and using a subsequence if necessary, that $\|\gamma_n\| < L \in \mathbb{R}$, for all $n \in \mathbb{N}$, for some constant L. From this we have that $(a(\gamma_n))$, $(b(\gamma_n))$, $(c(\gamma_n))$ and $(d(\gamma_n))$, are all bounded sequences with at least one of them non-constant, contradicting the discreteness of Γ. Hence (4) implies (3). Suppose that (3), and hence also (2), holds. If Γ were not discrete then there exists a sequence of distinct elements $(\gamma_n) \subset \Gamma$ and $\gamma \in \Gamma$ such that $\lim\limits_{n\to\infty} \gamma_n = \gamma$. But this would contradict the fact that $\lim\limits_{n\to\infty} \|\gamma_n\| = \infty$. This completes the proof. ∎

Lemma 4.1, extends a classical result due to Poincaré (see for example [12, Theorem II.1.2]). In particular, to check discreteness, one has to check discontinuity only at one point.

The proof of the lemma also shows that if $\Gamma < \text{PSL}(2,\mathbb{C})$ is discrete and $K \subset \mathbb{B}^3$ (respectively $\mathbb{B}^2$) is compact and (γ_n) is a sequence of distinct elements of Γ then there exists $n_0 \in \mathbb{N}$ such that $n \geq n_0$ implies that $K \cap \gamma_n(K) = \emptyset$. Since Γ is countable we have that $\{\gamma \in \Gamma \mid \gamma(K) \cap K\}$, is a finite subset of Γ.

Suppose that $\Gamma_0 = 1$ and put $r = r(\Gamma) = \min\{1 - \rho_\gamma \mid \gamma \in \Gamma\}$. Note that $r(\Gamma) = 1 - \rho_{\gamma_1}$, where $\|\gamma_1\| \leq \|\gamma\|$ for all $\gamma \in \Gamma - \{1\}$. By Lemma 3.5, the Euclidean ball $B_r(0)$ is contained in the fundamental domain of Γ with center 0. In particular, if Γ has finite covolume (respectively coarea) then $\text{vol}(\Gamma) \geq \pi(\sinh(2\rho(0,r)) - 2\rho(0,r))$ (respectively $\text{Area}(\Gamma) \geq 4\pi \sinh^2(\frac{1}{2}(\rho(0,r))))$. This can be used to give an estimate of $\text{vol}(\Gamma)$ (respectively $\text{Area}(\Gamma)$) for the examples in this paper. With the notation of Lemma 3.5, and using [12, Theorem II.5.4], there exists $\kappa \in]0,1[$ such that $\|\gamma\| \geq f^{-1}(1-\kappa)$, for all $\gamma \in \Gamma$ and Γ any discrete group. If $\Gamma_0 \neq 1$ we only have to consider a portion of the balls. This is the case in the first example of Section 4.2.

4.1 DF Domains

In this subsection we concentrate on discrete groups having a Poincaré fundamental domain which is also a Ford domain or which has two distinct points as center. Examples of such groups can be found in [33]. Here we revisit the interesting paper of [29] and complement its results. In particular we give an algebraic criterium that is easy to be checked. Our first result is the basis of what follows next.

Lemma 4.2 *Let $\gamma \in \text{PSL}(2,\mathbb{R})$. Then the following are equivalent.*

1. $a(\gamma) = d(\gamma)$
2. $\gamma = \sigma^2 \circ \tau \circ \sigma_\gamma$.
3. *Σ_γ is the bisector of the geodesic linking ti and $\gamma^{-1}(ti)$, for all $t > 0$.*

4. There exists $t_0 \neq 1$ such that Σ_γ is the bisector of the geodesic segment linking $t_0 i$ and $\gamma^{-1}(t_0 i)$.

Proof. Recall that $\sigma^2 \circ \tau$ denotes the reflection in the imaginary axes, i.e., $\sigma^2 \circ \tau(z) = -\overline{z}$. We first prove that the two first items are equivalent.

Suppose that $a(\gamma) = d(\gamma)$. Then, by Proposition 3.3, we have that $\Sigma_\gamma = \mathrm{ISO}_\gamma$. Hence the reflection σ_γ in Σ_γ is given by $\sigma_\gamma(z) = P_\gamma - (|c|^2 \sigma^2 \circ \tau(z - P_\gamma))^{-1} = \sigma^2 \circ \tau(\gamma(z))$, if $c(\gamma) \neq 0$. If $c(\gamma) = 0$, then we may take $a(\gamma) = d(\gamma) = 1$ and thus $\sigma_\gamma(z) = \sigma^2 \circ \tau(z - b(\gamma)) = \sigma^2 \circ \tau(\gamma(z))$. Hence in either case we have that $\gamma = \sigma^2 \circ \tau \circ \sigma_\gamma$. Now suppose that $\gamma = \sigma^2 \circ \tau \circ \sigma_\gamma$. We first suppose that P_γ exists, i.e., Σ_γ is an Euclidean sphere. In this case we have that $\gamma(z) = \sigma^2 \circ \tau \circ \sigma_\gamma(z)$ $= \frac{-P_\gamma z + P_\gamma^2 - R_\gamma^2}{z - P_\gamma}$, from which it follows that $a(\gamma) = d(\gamma)$. If Σ_γ is a vertical line, $x = x_0$ say, then $\sigma_\gamma(z) = -\overline{z} + 2x_0$ Hence $\gamma(z) = \sigma^2 \circ \tau \circ \sigma_\gamma(z) = z - 2x_0$ and hence $a(\gamma) = d(\gamma)$.

Let $u \in \Sigma_\gamma$. Then $\rho(u, \gamma^{-1}(ti)) = \rho(\gamma^{-1}(ti), u) = \rho(\sigma_\gamma \circ \sigma^2 \circ \tau(ti), u) = \rho(\sigma_\gamma(ti), u) = \rho(ti, \sigma_\gamma(u)) = \rho(ti, u)$ and hence Σ_γ is the bisector of the geodesic linking ti and $\gamma^{-1}(ti)$. This proves that the first item implies the third. Obviously the third item implies the fourth.

We now prove that the fourth item implies the first. Let $u \in \Sigma_\gamma$. Then we have that $\rho(ti, u) = \rho(u, \gamma^{-1}(ti))$ and hence $\rho(ti, u) = \rho(ti, \gamma(u))$. Since γ is a Möbius transformation we have that $\mathrm{Im}(\gamma(z)) = |\gamma'(z)| \mathrm{Im}(z)$. From this we obtain that $|\gamma\prime(u)||ti - u|^2 = |ti - \gamma(u)|^2$. It follows that $\mathrm{Re}(u)^2 - \mathrm{Re}(\gamma(u))^2 |\gamma'(u)| = (|\gamma'(u)| - 1)\mathrm{Im}(u)t^2 + (1 - |\gamma'(u)|)|\gamma'(u)|\mathrm{Im}(u)^2$. We may write this as an equation of the type $at^2 = c$ having $t = 1$ and $t = t_0$ as solutions and hence $a(t_0^2 - 1) = 0$. It follows that $a = 0$ and thus $|\gamma'(u)| = 1$, for all $u \in \Sigma_\gamma$, i.e., $\Sigma_\gamma = \mathrm{ISO}_\gamma$. Applying Proposition 3.3, we obtain that $a(\gamma) = d(\gamma)$. ∎

The following corollary is useful to decide if a set of matrices generates a discrete group. For example it can be used to prove that the group Γ of [29, Section VI] is discrete.

Corollary 4.3 *Let $\{\gamma_1, \cdots, \gamma_n\} \subset \mathrm{PSL}(2, \mathbb{R})$ and suppose that $a(\gamma_k) = d(\gamma_k)$ for all $1 \leq k \leq n$. Then $\langle \gamma_1, \cdots, \gamma_n \rangle$ is Fuchsian if and only if all dihedral angles of intersecting isometric spheres of the γ_k's are submultiples of π. In particular, $\langle \gamma_1, \cdots, \gamma_n \rangle$ is the subgroup of the orientation preserving isometries of a Fuchsian reflection group.*

Proof. We only have to prove the converse. If Σ_{γ_k} intersects the imaginary axis Σ, say, then so does $\Sigma_{\gamma_k^{-1}}$ and $(\Sigma, \Sigma_{\gamma_k}) = \frac{1}{2}(\Sigma_{\gamma_k}, \Sigma_{\gamma_k^{-1}})$. In the elliptic case $(\Sigma, \Sigma_{\gamma_k}) = \frac{\pi}{2}$. In particular, $(\Sigma, \Sigma_{\gamma_k})$ is a submultiple of π. Consider the polyhedron P, say, whose sides are Σ and the Σ_{γ_k}'s with $P_{\gamma_k} \geq 0$. By [33, Theorem VII.1.3], the group $\tilde{\Gamma}$ generated by the reflections of $\mathbb{H}^2$ in the sides of P is a discrete reflection group with respect to P. By Lemma 4.2 we have that $\tilde{\Gamma} = \langle \sigma^2 \circ \tau, \sigma_{\gamma_k} \mid P_{\gamma_k} \geq 0 \rangle$ is a reflection group containing $\langle \gamma_1, \cdots, \gamma_n \rangle$. Since $\langle \gamma_1, \cdots, \gamma_n \rangle = \langle \gamma_k \mid P_{\gamma_k} \geq 0 \rangle$, we have that $\langle \gamma_1, \cdots, \gamma_n \rangle$ is of index two in the reflection group $\tilde{\Gamma}$. ∎

In [29, Theorem 5.3], conditions are given for a Fuchsian group Γ to have a fundamental domain $\mathcal{F}$ which is both a Ford domain and a Dirichlet domain (a DF domain): *A finitely generated, finite coarea Fuchsian group Γ admits a DF domain $\mathcal{F}$, if and only if Γ is an index 2 subgroup of a reflection group.* It is also proved that in the Kleinian case Γ has a generating set consisting of elements whose traces are real ([29, Theorem 6.3].) Our next theorem complements this nice result.

Theorem 4.4 *Let $\Gamma < \mathrm{PSL}(2, \mathbb{C})$ be a finitely generated discrete group, acting on $\mathbb{H}^n, n \in \{2, 3\}$, and $P_0 \in \{i, j\}$, according to Γ being Fuchsian or Kleinian. Then Γ admits a DF domain with* *center* P_0 *if and only if for every side-pairing transformation* γ *of* $\mathcal{F}$ *we have that* $d(\gamma) = \overline{a(\gamma)}$. *In particular,* $\mathrm{tr}(\gamma) \in \mathbb{R}$ *for all these elements. Moreover, if* Γ *is Fuchsian, then* $\tilde{\Gamma} = \langle \sigma^2 \circ \tau, \Gamma \rangle$ *is a reflection group,* $\hat{\Gamma} = \langle \sigma^2, \Gamma \rangle$ *is a Coxeter Kleinian group and both contain* Γ *as a subgroup of index two.*

Proof. Let $\mathcal{F}$ be a DF domain, in $\mathbb{H}^3$, for Γ with center $P_0 \in \{i, j\}$. Let Φ be the side-pairing transformations of $\mathcal{F}$ and let Φ_0 consist of those elements of Γ whose isometric circles (respectively isometric spheres) and vertical lines (respectively vertical planes) form the boundary of $\mathcal{F}$.

Then there exists a bijection $f : \Phi_0 \ \to \Phi$ such that if $\gamma \in \Phi_0, \gamma \notin \Gamma_\infty$ then $\mathrm{ISO}_\gamma = \Sigma_{f(\gamma)}$. Since $\mathcal{F}$ is a Ford domain we have that $\mathcal{F} \cap \gamma^{-1}(\mathcal{F}) = \mathrm{ISO}_\gamma$ and hence $\gamma \in \Phi$. Since $\mathcal{F}$ is also a Dirichlet domain we have that $\mathcal{F} \cap \gamma^{-1}(\mathcal{F}) = \Sigma_\gamma$. Consequently $\mathrm{ISO}_\gamma = \Sigma_\gamma$ and thus, by Proposition 3.3, $d(\gamma) = \overline{a(\gamma)}$. It also follows that $\Sigma_\gamma = \Sigma_{f(\gamma)}$ and hence, by item 7 in Proposition 3.4, $f(\gamma) = g\gamma$, with $g \in \Gamma_{P_0}$.

If Σ_γ is an Euclidean line or plane then, since f is a bijection, $\Sigma_{\gamma^{-1}}$ is also a line or a plane. From this it follows that $|a(\gamma)| = |d(\gamma)| = 1 = a(\gamma)d(\gamma)$ and thus $d(\gamma) = \overline{a(\gamma)}$ and $c(\gamma) = 0$. Hence, in the Kleinian case, $\mathrm{tr}(\gamma) \in \mathbb{R}$, for all $\gamma \in \Phi$.

To prove the converse, one just has to use Proposition 3.3 to obtain that $\mathrm{ISO}_\gamma = \Sigma_\gamma$, for all side paring transformation γ whose bisector is not an Euclidean line or plane, and hence we have a DF domain.

Suppose now that Γ is Fuchsian. In this case, the set of elements of Φ_0 without an isometric circle is $\Phi_0 \cap \Gamma_\infty$ and consists of only two elements, γ_0 and γ_0^{-1} say. Letting $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \gamma = f(\gamma_0)$, we have that Σ_γ is an Euclidean line, $|a|^2 + |c|^2 = 1$, and $x = -\frac{ab+cd}{2}$ is a defining equation of Σ_γ. Since Γ has finite coarea, γ_0 is parabolic and hence $\gamma_0 = \begin{pmatrix} 1 & -b_0 \\ 0 & 1 \end{pmatrix}$. Clearly $f(\gamma_0^{-1}) = f(\gamma_0)^{-1}$, hence $|d|^2 + |c|^2 = 1$, and thus $|a|^2 = |d|^2$. Suppose that $c \neq 0$; we have that $\infty \in \Sigma_\gamma$ and $\gamma(\infty) = \frac{a}{c}$. Since $\gamma(\Sigma_\gamma) = \Sigma_{\gamma^{-1}}$, it follows that $\frac{a}{c} = \frac{db+ac}{2}$. If $a = 0$ then $\gamma_0 \in \Gamma_i$. If $a \neq 0$ then $d = -a$. We obtain that $a^2 + bc = -1$ and hence $\gamma^2 = -1$. Hence $o(\gamma) = 2$ and thus $\Sigma_\gamma = \Sigma_{\gamma^{-1}}$, a contradiction. So we must have that $c = 0$, hence $a^2 = d^2 = 1 = ad$, $a = d = 1$ and $b = b_0$. This proves that for any $\gamma \in \Phi$ we have that $d(\gamma) = a(\gamma)$.

By Corollary 4.3 we have that $\tilde{\Gamma} = \langle \sigma^2 \circ \tau, \Phi \rangle$ is a reflection group containing Γ as a subgroup of index 2. Since $o(\sigma^2\gamma) = 2$, $\mathrm{tr}(\sigma^2\gamma) = 0$, for all $\gamma \in \Phi$ it follows that $\hat{\Gamma} := \langle \sigma^2, \Phi \rangle$ is a Coxeter group with $[\hat{\Gamma} : \Gamma] = 2$. ▮

Note that a presentation of $\tilde{\Gamma}$ and $\hat{\Gamma}$ can be obtained using [12, Theorem II.7.5]. In the Fuchsian case it easily follows that the orbifold of Γ is a punctured sphere (see [29]).

Corollary 4.5 *Let* Γ *be a Fuchsian group. Then the following are equivalent.*

1. Γ *is the subgroup of orientation preserving isometries of a Fuchsian reflection group.*
2. Γ *has a DF domain.*
3. Γ *has a Poincaré Fundamental domain* $\mathcal{F}$ *such that* $a(\gamma) = d(\gamma)$ *for every side-pairing transformation* γ.
4. Γ *has a Poincaré Fundamental domain* $\mathcal{F}$ *with two distinct points as center.*

5. Γ *has a Poincaré Fundamental domain* $\mathcal{F}$ *with a geodesic ray all whose points are centers of* $\mathcal{F}$.

Proof. We just have to prove that the first item implies the second. Fix a polyhedron P for the reflection group such that one of the sides is the imaginary axis Σ, say. Then $\sigma^2 \circ \tau$ is the reflection in Σ. Let Σ_i be a side of P, denote by σ_{γ_i} the reflection in Σ_i and let $\gamma_i = \sigma^2 \circ \tau \circ \sigma_{\gamma_i}$. Then clearly $\Sigma_i = \Sigma_{\gamma_i}$ and the rest follows by Lemma 4.2. ∎

Our proof permits that Γ_j is non-trivial. In this case $S^n \cap \mathbb{H}$ is the isometric sphere of $\gamma(P) = -P^{-1}$ and this can be part of the boundary of a fundamental domain of Γ_{P_0}. It follows that all examples of [12, Section VII.4] are groups whose Ford domain is also a Dirichlet domain. Note that this does not follow from the results of [29].

Another interesting example is the group inducing the unique hyperbolic structure of the trice-punctured sphere. This is the free group on two generators $\Gamma(2) = \langle z+2, \frac{z}{2z+1} \rangle$, the congruence group at level 2. Its orbifold is the trice-punctured sphere (see for example [33, Theorem 9.8.8]). By the previous theorem it has a DF domain. The same can be deduced for $\mathrm{PSL}(2,\mathbb{Z})$. Hence we see that $\langle \Gamma(2), \sigma^2 \circ \tau \rangle < \langle \mathrm{PSL}(2,\mathbb{Z}), \sigma^2 \circ \tau \rangle$ are both reflection groups and hence $\langle \Gamma(2), \sigma^2 \circ \tau \rangle$ is not a maximal reflection group. In this case this is due to one more symmetry in the fundamental domain of $\Gamma(2)$. In [29, Section VI] a maximal reflection group, constructed in this way, is exhibited thus answering a question of [1]. The fundamental domain does not have the symmetric property exhibited by the fundamental domain of $\Gamma(2)$. Note that only visual symmetry is not enough.

Before continuing the investigations, we describe a general procedure on how to use the results of the previous section to construct a fundamental domain of a group $\Gamma \subset \mathrm{Iso}^+(X)$ that is discrete, of finite covolume and X is the hyperbolic two or three space. We concentrate on describing how to obtain generators for a subgroup of finite index of Γ. Using, Lemma 3.5, the procedure can be refined to find generators for Γ. This is explained in the next section and executed for d=15 (Theorem 4.8). We shall work in $\mathbb{B}^n$ to make things clearer. We use η_0 to switch between the two models. We now describe the *Dirichlet Algorithm of Finite Covolume* for $n \in \{2,3\}$.

The DAFC

I Embed Γ in $\mathrm{PSL}(2,\mathbb{C})$.
II Find a system of algebraic equations (X) whose set of solutions is Γ.
III Take $f : \Gamma \to \mathbb{R}$, where $f(\gamma) = \|\gamma\|^2$.
IV Take an increasing sequence (r_n) such that $(r_n) = \mathrm{Im}(f)$.
V For each $n \in \mathbb{N}$ draw all the bisectors of the elements of the set $f^{-1}(r_n)$.
VI Stop when $\partial\mathbb{B}^n \subset \bigcup_{f(\gamma) \le r_{n_0}} \overline{B}_{R_{\Psi(\gamma)}}(P_{\Psi(\gamma)})$, for some $n_0 \in \mathbb{N}$.
VII Use η_0^{-1} to obtain a fundamental domain in $\mathbb{H}^n$ of a subgroup of finite index.

Note that $f^{-1}(x)$ is always finite because Γ is discrete. This is well known and can also be seen as follows: If $\|\gamma_1\| = \|\gamma_2\|$ then, by Proposition 3.4, $\|P^*_{\Psi(\gamma_1)}\| = \|P^*_{\Psi(\gamma_2)}\|$. Since $P^*_{\Psi(\gamma)} = \Psi(\gamma^{-1})(0) \in \mathbb{B}^3$ and Γ is discrete, the claim follows. Since Γ is of finite covolume, results of Greenberg, Garland and Raghunathan (see [12, Theorem II.2.7]) imply that the DAFC stops in a finite number of steps. In fact, it stops in a finite number of steps for any geometrically finite group. In the next two subsections and in Section 5, we show how to implement the DAFC.

In case we have $\Gamma = \langle X \rangle$ we can still make use of DAFC. In this case we set $S = X \cup X^{-1}$, where $X^{-1} = \{x^{-1} \in X \mid x \in X\}$ and make the Cayley graph $\mathcal{G}(\Gamma)$ of Γ using S. The word length metric of $\mathcal{G}(\Gamma)$ gives a natural way of turning Γ into a sequence of elements. Now use this sequence in item IV of DAFC.

4.2 Division Algebras

In this subsection all examples given are discrete and cocompact subgroups of $\mathrm{Iso}^{+}(\mathbb{B}^n), n = 2, 3$. One encounters two situations.

Case I: Γ_0 is non-trivial. In this case we first determine a fundamental domain $\mathcal{F}_0$ of Γ_0, a polyhedron with the origin on its boundary, and let $V = \partial(\mathcal{F}_0) \cap \partial\mathbb{B}^n$. Using the DAFC, find a finite set $\{\gamma_1, \cdots, \gamma_n\} \subset \Gamma$, such that $V \subset \bigcup\limits_{1 \le k \le n} \mathrm{Interior}(\Sigma_{\Psi(\gamma_k)})$. As output we get $\langle \Gamma_0, \gamma_1, \cdots, \gamma_n \rangle$ which has finite index in Γ.

Case II: Γ_0 is trivial. In this case we first determine $\gamma_0 \in \Gamma$ whose bisector Σ separates j and the origin 0. Because of Lemma 3.6, the existence of such a γ_0 is guaranteed and furthermore $\gamma_0 = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $|a|^2 + |c|^2 < 1$. Let $V = \mathrm{Exterior}(\Sigma_{\Psi(\gamma_0)}) \cap S^2$ and proceed as in case I.

Generators for Γ: Once obtained a fundamental domain $\mathcal{F}_1$, say, of a subgroup of finite index in Γ, one finds a set of generators, and hence a fundamental domain for Γ, as follows. Let $t = \max\{\|v\| \mid v \in \partial\mathcal{F}_1\}$ and find $n_0 \in \mathbb{N}$ such that $m > n + n_0$ implies that $\rho_{\gamma_m} < 1 - t$. Then $\Gamma = \langle \Gamma_0, \gamma_1, \cdots, \gamma_n, \cdots \gamma_{n+n_0} \rangle$. Note that t can be calculated looking only at the vertices of $\mathcal{F}_1$.

One observation with regard to **Case II**: in all the examples in this subsection we will have $c = 0$, which makes this search easier since this amounts to finding a unit in the ring of algebraic integers of a maximal field of the algebra involved (see the last three examples of this subsection).

We now have all the tools to implement the DAFC in a way that the reader can follow and easily reproduce.

Our first example is a generalized quaternion division algebra $\mathcal{H}(a, b, K)$ with $K = \mathbb{Q}(\sqrt{-d}), d \in \mathbb{N}$. In this case, the unit group of any order of $\mathcal{H}(K)$ is a cocompact Kleinian group (see [12, Theorem X.1.2]). In particular, we consider Example 8 of chapter X of [12] (page 474), one example of [26] and also revisit the work of [10].

To implement the DAFC, we find an additive basis for the ring of integers of K. This, together with the fact that we only consider reduced norm one elements in the unit group, leads to a system of Diophantine equations whose solution set is Γ. We get a sieve (the function f), parameterizing the system by the matrix norm of the elements. For small values of d the algorithm easily can be done by hand, and for larger d, we made use of the software package Mathematica, to get a Poincaré fundamental domain and hence a set of generators (up to finite index).

We first revisit [10]. In this case we get a particular nice system of Diophantine equations one of which is to write a number as the sum of four squares.

Consider $\mathcal{H}(K)$, with $K = \mathbb{Q}(\sqrt{-d})$, with d a positive square free integer. We will consider the cases $d = 15$ and $d = 23$. Suppose that $d \equiv 3 \bmod 4$. When $d \equiv 7 \bmod 8$ then $\mathcal{H}(K)$ is a division algebra and thus $\Gamma = SL_1(\mathcal{H}(-1, -1, \mathfrak{o}_K)) = \{x \in \mathcal{H}(-1, -1, \mathfrak{o}_K) : N(x) = 1\}$, acts discretely and cocompactly on $\mathbb{H}^3$. In all other cases $\mathcal{H}(K)$ is a matrix algebra and hence, since its elements satisfy the hypothesis of Corollary 3.7, it follows that in these cases Γ is never cocompact (see also Chapter VII of [12]).

It is easily seen that $\Gamma_0 = \langle i, j\rangle \cong Q_8$, the quaternion group of order 8 and that $\eta_0^{-1}(\mathcal{F}_0)$ can be taken to be that part of the unit ball centered at the origin whose projection on $\partial\mathbb{H}^3$ is the upper half of the unit circle, $\{z \in \mathbb{C} \mid |z| < 1, Im(z) > 0\}$.

We have that $\mathfrak{o}_K = \text{span}_{\mathbb{Z}}[1, w]$, where $w = \frac{1+\sqrt{-d}}{2}$. Write $u = u_0 + u_1 i + u_2 j + u_3 k \in \mathcal{H}(K)$ and $u_t = x_t + y_t w$. Define $x = (x_0, x_1, x_2, x_3)$ and $y = (y_0, y_1, y_2, y_3)$. So u is determined by the vector (x, y). We will use this notation to identify units. Furthermore, in this example we use the embedding $u \mapsto \gamma_u = \begin{pmatrix} u_0 + u_1 i & u_2 + u_3 i \\ -u_2 + u_3 i & u_0 - u_1 i \end{pmatrix}$ as used in [10].

The next lemma gives formulas for Step II of the DAFC.

Lemma 4.6 *Let $u = u_0 + u_1 i + u_2 j + u_3 k \in \mathcal{H}(K)$, $\gamma = \gamma_u$ and $N(u) = \pm 1$. Then*

$$\begin{cases} \|x\|^2 - (\frac{d+1}{4})\|y\|^2 = N(u) \\ 2\langle x|y\rangle + \|y\|^2 = 0 \end{cases} \tag{1}$$

Moreover, the following hold.

1. $\|\gamma_u\|^2 = 2N(u) + d\|y\|^2 \in 2\mathbb{Z}$
2. $\|P_{\Psi(\gamma)}\|^2 = \frac{2+2N(u)+d\|y\|^2}{-2+2N(u)+d\|y\|^2}$
3. $R^2_{\Psi(\gamma)} = \frac{4}{-2+2N(u)+d\|y\|^2}$.

Proof. The first part of the statement follows by using the integral basis and the condition on the reduced norm. The second part follows from Proposition 3.4. ▮

We shall say that (x, y) satisfies a Pell-Vector-Equation (PVE) if (x, y) satisfies the first set of equations of the previous lemma. If we set $\|y\|^2 = 2n \in \mathbb{N}$ then we are in position to apply the DAFC.

Let $J(x) = (x_1, -x_0, -x_3, x_2)$ and $S(x) = (-x_3, x_2, -x_1, x_0)$. Then J and S are skew orthogonal linear maps and $\langle S, J\rangle$ is isomorphic to Q_8.

Lemma 4.7 *Let $u \in \mathcal{H}(K), N(u) = \pm 1$ and $\gamma = \gamma_u = \begin{pmatrix} u_0 + u_1 i & u_2 + u_3 i \\ -u_2 + u_3 i & u_0 - u_1 i \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then*

1. $|a|^2 + |c|^2 = N(u) + \frac{d}{2}\|y\|^2 + \langle J(x)|y\rangle \cdot \sqrt{d}$ *and* $|b|^2 + |d|^2 = N(u) + \frac{d}{2}\|y\|^2 - \langle J(x)|y\rangle \cdot \sqrt{d}$
2. $\overline{a}b + \overline{c}d = [\langle -S(x)|y\rangle + i\langle SJ(x)|y\rangle] \cdot \sqrt{d}$
3. $|\overline{a}b + \overline{c}d|^2 = (N(u) + \frac{d}{2}\|y\|^2)^2 - d\langle J(x)|y\rangle^2 - N(u)$
4. *If* $N(u) = 1$ *then* $R_{\Psi(\gamma)} = \frac{2\sqrt{d}}{d\|y\|}$
5. *If* $N(u) = 1$ *then* $P_{\Psi(\gamma)} = \frac{2\sqrt{d}}{d\|y\|^2} \cdot [\langle S(x)|y\rangle - \langle S(x)|J(y)\rangle i - \langle J(x)|y\rangle j\ ]$

The formulas show that the centers of the bisectors in the ball model, up to a scalar in $\mathbb{Z}[\sqrt{d}]$, belong to $\mathbb{Z}^3$ and in the upper half space model they belong to $\mathbb{Z}[i]$. The group of symmetries of a fundamental domain of Γ contains $\langle\sigma,\tau\rangle$ and $\langle\Gamma,\sigma,\tau\rangle$ is a discrete, probably maximal, group. Note that ϕ also acts on the tessellation of $\mathbb{H}^3$ induced by Γ. However since $S^2\cap\mathbb{H}^3$ is part of the boundary of Γ_j, this symmetry is lost. It shows up if we work with a torsion free subgroup of Γ.

Theorem 4.8 *Let* $\Gamma=\mathrm{SL}_1(\mathcal{H}(-1,\,-1,\mathbb{Z}[\frac{1+\sqrt{-d}}{2}]))$ *and let* X_d *be a finite set of units whose bisectors cover the circular sector* $\{(a,b)\mid a^2+b^2\le 1,\ 0\le b\le a\}$ *and let* $S_d=\Gamma_0\cup\{g(\gamma)\mid g\in\langle\sigma,\tau\rangle,\ g\in X_d\}$.

1. *If* $d=15$ *then* $\Gamma=\langle S_d\rangle$*, where*

$$\begin{aligned}X_{15}\quad=\quad&\{(2,-1,2,0,0,1,0,1),(2,2,1,0,0,0,-1,1),(-2,-2,1,0,0,1,1,0),\\&(-2,-1,0,-2,0,1,1,0),(-2,1,2,0,1,1,0,0),(-4,2,3,-2,2,0,0,-2),\\&(-4,3,2,-2,2,0,0,-2),(-4,3,-2,-2,2,0,0,-2),(-4,2,3,-2,2,2,0,0)\}\,.\end{aligned}$$

2. *If* $d=23$ *then* $\langle S_d\rangle$ *has finite index in* Γ*, where*

$$\begin{aligned}X_{23}\quad=\quad&\{(-3,2,0,0,1,1,0,0),(-2,-2,1,-2,1,0,1,0),(-3,0,0,-2,1,0,0,-1),\\&(-2,-2,-2,-1,0,0,1,-1),(-2,-2,1,-2,0,1,1,0),(-4,2,-2,1,0,0,0,-2),\\&(-4,-1,2,2,0,2,0,0),(-4,-2,1,-4,2,0,1,-1),(-5,2,-2,-2,1,-1,0,-2),\\&(-4,1,2,-4,2,1,0,-1),(-4,1,-2,4,-1,1,0,-2),(-5,2,-2,-2,1,0,1,-2),\\&(-4,-4,1,2,-1,2,1,0),(-5,2,2,2,1,2,-1,0),(-6,-3,-2,0,0,0,2,-2),\\&(-5,2,2,4,0,2,0,-2),(-6,0,3,2,0,2,0,-2),(-5,-2,-4,-2,0,0,2,-2),\\&(-5,-4,2,-2,0,2,2,0),(-6,0,-3,2,0,2,0,-2)\}\,.\end{aligned}$$

Part (a) of the figures below shows the projection on $\partial\mathbb{H}^3$ of the bisectors of the elements in X_d and part (b) shows the fundamental domain of the subgroup $\langle S_d\rangle$.



(a) Projection of fundamental domain of $\langle S_{15}\rangle$.

(b) Fundamental domain of $\langle S_{15}\rangle$.

Figure 1:



(a) Projection of fundamental domain of $\langle S_{23}\rangle$.

(b) Fundamental domain of $\langle S_{23}\rangle$.

Figure 2:

For each d we have that $r = r(\Gamma) = \sqrt{\frac{2+d}{d}} - \sqrt{\frac{2}{d}}$. Since $\Gamma_0 \neq 1$, we get that $\mathrm{vol}(\Gamma) \geq \frac{\pi}{4}(\sinh(2\rho(0,r)) - 2\rho(0,r))$. It follows that $\lim\limits_{d\to\infty} \mathrm{vol}(\Gamma) = \infty$ and the convergence is exponential.

Although we used Mathematica, all calculations can be done quickly by hand for $d \in \{7, 15, 23\}$. To do so we first solve the sum of four squares equations $\|y\|^2 = 2n, n \in \mathbb{N}$, next solve the sum of four squares equations $\|x\|^2 = 1 + (\frac{d+1}{4})\|y\|^2$ and check if $2\langle x|y\rangle + \|y\|^2 = 0$. In this way, we find all units beginning with those of norm 2. We may concentrate on solutions y, all whose coordinates are non-negative, leaving the negative sign, if necessary, for the coordinates of x. Let G_0 be the subgroup of $\mathrm{SL}(4,\mathbb{Z})$ generated by the orthogonal matrices whose columns are in $\{\pm e_i \mid i = 1,2,3,4\}$, where the e_i's are the elements of the canonical bases of $\mathbb{R}^4$. If $g \in G_0$ and (x,y) defines a unit then (gx, gy) also defines a unit. Using this, and the symmetries defined above, the task of finding a subgroup of finite index is now easy. As an example take $d = 7$. Then $\|y\|^2 = 2n, \|\gamma\|^2 = 2 + 2dn = 2 + 14n, n \in \mathbb{N}$. So $\|\gamma\|^2$ is an arithmetic progression. If $n = 0$ we have the trivial units $\{\pm i, \pm j, \pm k\}$. If $n = 1$, making use of the action of G_0, we may take $y = (1,1,0,0)$. We have $\|x\|^2 = 9, x_0 + x_1 = -1$ which gives us the following solutions: $x \in \{\ (0,-1,\pm 2,\pm 2)$, $(1,-2,\pm -2,0)$, $(1,0,\pm 2,\pm 2)$, $(-2,1,\pm 2,0)\}$. Some of the units are $u_1 = (-2+w) + (1+w)i + 2j$, $u_2 = -1 + w + wi + 2j + 2k$, $u_3 = 1 + w + (-2+w) + 2j$ and $u_4 = w + (-1+w)i + 2j + 2k$. Note that if Σ_{γ_u} does not intersect the unit ball then $\Sigma_{\gamma_{uj}}$ does intersect it. In fact, $\gamma_{uj} = \delta(\gamma_u)$ and thus $P_{\gamma_{uj}} = P_{\delta(\gamma_u)}$. Using these units and the symmetries, we can now quickly cover the circular sector mentioned in the previous theorem. One important consequence is that we can give an explicit description of all units of this order, for all d.

In the context of a division algebra, Pell and Gauss units were constructed in [24, 25]. Those units for which $P_\gamma = 0$, are typically associated to hyperbolic elements. They also arise from a Pell equation. Together, these units seem to generate a subgroup of finite index. If so, these can be used as generic units in the group ring problem. Still in the same context, in [24] it was proved that $SL_1(\mathcal{H}(-1,-1,\mathfrak{o}_K))$ is a hyperbolic group for all d. This line of classification was introduced and initiated in [23].

The next cases we treat are division algebras of the form $\mathcal{H}(a,b,\mathbb{Q}(i'))$, $(i' = \sqrt{-1})$, with $0 < a < b$ integers and $\Gamma = \mathrm{SL}_1(\mathcal{H}(a,b,\mathbb{Z}[i']))$. For $a = 2$ and $b = 5$, this can be found in [12, Section

X], and it is commensurable with $\mathrm{PO}_4(\mathbb{Z},q)$, where $q(x,y,z,w)=-2x^2-5y^2-10z^2+w^2$.

Let $u\in\Gamma$ and write $u=u_0+u_1i+u_2j+u_3k$, with $u_t\in\mathbb{Z}[i']$. In this case $\gamma_u=\begin{pmatrix}u_0+u_1\sqrt{a} & u_2\sqrt{b}+u_3\sqrt{ab}\\ u_2\sqrt{b}-u_3\sqrt{ab} & u_0-u_1\sqrt{a}\end{pmatrix}$. Let $u_t=x_t+y_ti'$, $x=(x_0,x_1,x_2,x_3)$, $y=(y_0,y_1,y_2,y_3)$, $q(x)=x_0^2-ax_1^2-bx_2^2-abx_3^2$ and let $B(x,y)$ be the bilinear form associated to $q(x)$. Then we obtain the following system of Diophantine equations to describe Γ as a subgroup of $\mathrm{PSL}(2,\mathbb{C})$..

$$\begin{cases}q(x)-q(y)=1\\ B(x,y)=0\end{cases}$$

Letting $x_o^2+ay_1^2+by_2^2+abx_3^2=n\in\mathbb{N}$, we obtain

$$\begin{cases}x_o^2+ay_1^2+by_2^2+abx_3^2=n\\ y_0^2+ax_1^2+bx_2^2+aby_3^2=n-1\\ B(x,y)=0\\ \|\gamma_u\|^2=4n-2\end{cases}\tag{2}$$

Here $\Gamma_j=1$ and so we are in Case II. Taking $a=2,b=5$ and $n=2$ in system (2), we find u such that $\gamma_u=\begin{pmatrix}i'-i'\sqrt{2} & 0\\ 0 & i'+i'\sqrt{2}\end{pmatrix}$ and thus $|a|^2+|c|^2=3-2\sqrt{2}<1$. Using Mathematica, we find the needed units, all whose norm are less then $4\cdot 102-2$. In this case $\langle\sigma,\tau,\phi\rangle$ is contained in the group of symmetries of Γ.

Theorem 4.9 *In* $\mathrm{SL}_1(\mathcal{H}(2,5,\mathbb{Z}[i']))$, *the subgroup* $S_{2,5}(\mathbb{Z}[i])=\langle -1,g(\gamma)\mid g\in\langle\sigma^2,\tau,\phi\rangle,\gamma\in X\rangle$, *where* X *is given below, has finite index.*

$$\begin{aligned}X\;=\;&\{i'-i'i,2i'-i'j,2-i'i-j,2-2i+i'j,3i'+k,3+i'i-i'k,2+i'i-i'j-i'k,\\ &4i'-i'i-i'j+k,3-3i+k,2-3i+i'j+k,3-i'i-2j-k,6-i'j-2i'k,\\ &2+4i'i-i'j-2i'k,6-3j-k,2-4i'i-3j-k,(1-4i')+(4+2i')i+(2+2i')j+2k,\\ &(1+4i')+(4-2i')i+(-2-2i')j-2k,3-7i+4i'j+k,\\ &(8+3i')+(-4+2i')i+(-2-2i')j+(1-2i')k\}\,.\end{aligned}$$

Figures 3(a) and 3(b) show a part of the fundamental domain and the projection on $\partial\mathbb{H}^3$ of the fundamental domain. The bisectors coming from the elements in X are drawn in bold face.



(a) Projection of the fundamental domain of $S_{2,5}(\mathbb{Z}[i])$. (b) Part of the fundamental domain of $S_{2,5}(\mathbb{Z}[i])$.

Figure 3:

In this case $r(\Gamma) \geq \sqrt{\frac{a}{a-1}} - \sqrt{\frac{1}{a-1}}$ and with this we get an estimate for the volume of Γ.

We finish this section with two examples of orders in division algebras over $\mathbb{Q}$, i.e., with two Fuchsian groups. The first example is taken from [26]. Let $a, b \in \mathbb{N}$ and $\Gamma_{ab} < \mathrm{PSL}(2,\mathbb{R})$ be given by

$$\Gamma_{ab} = \{\gamma = \begin{pmatrix} \frac{l+m\sqrt{a}}{2} & \sqrt{b}(\frac{w-u\sqrt{a}}{2}) \\ \sqrt{b}(\frac{w+u\sqrt{a}}{2}) & \frac{l-m\sqrt{a}}{2} \end{pmatrix}, (l,m,u,w) \in \mathbb{Z}^4, \\ l \equiv w (\mathrm{mod}\ 2), m \equiv u (\mathrm{mod}\ 2), \det(\gamma) = 1\}$$

The condition on the norm gives us that

$$\begin{cases} l^2 + abu^2 = n \\ am^2 + bw^2 = n - 4 \\ l \equiv w (\mathrm{mod}\ 2), m \equiv u (\mathrm{mod}\ 2), \end{cases} \tag{3}$$

$\|\gamma\|^2 = n - 2$, $n \in \mathbb{N}, n \geq 4$. Supposing that $a = 3$ and $b = 5$ we get for $n = 4$ the identity and hence the stabilizer $\Gamma_i = 1$. Taking $n = 16$, we find $\gamma_u = \begin{pmatrix} 2-\sqrt{3} & 0 \\ 0 & 2+\sqrt{3} \end{pmatrix}$, such that $|a|^2 + |c|^2 = 7 - 4\sqrt{3} < 1$. Using the solutions for $n = 9$ and $n = 31$, we obtain generators for a subgroup of finite index. In this case, it is easy to see that this is a fundamental domain of the whole group. Here the symmetry reduces to just one, namely, σ^2.

Theorem 4.10 *The group* Γ_{15} *is generated by the set*
$$\left\{\begin{pmatrix}\frac{3}{2} & \frac{\sqrt{5}}{2}\\ \frac{\sqrt{5}}{2} & \frac{3}{2}\end{pmatrix}, \begin{pmatrix}2+\sqrt{3} & 0\\ 0 & 2-\sqrt{3}\end{pmatrix}, \begin{pmatrix}\frac{4+3\sqrt{3}}{2} & -\frac{\sqrt{15}}{2}\\ \frac{\sqrt{15}}{2} & \frac{4-3\sqrt{3}}{2}\end{pmatrix}, \begin{pmatrix}\frac{4-3\sqrt{3}}{2} & -\frac{\sqrt{15}}{2}\\ \frac{\sqrt{15}}{2} & \frac{4+3\sqrt{3}}{2}\end{pmatrix}\right\}$$



Figure 4: Fundamental domain of Γ_{15}

In this last case $r(\Gamma) = \frac{3\sqrt{5}}{5} - \sqrt{\frac{4}{5}}$ and from this we can obtain an estimate of the coarea of Γ.

Finally, consider $\mathcal{H}(a,b,\mathbb{Z}) \subset \mathcal{H}(a,b,\mathbb{Q})$ with $a>0, b>0$ integers. For $u = x_0 + x_1 i + x_2 j + x_3 k \in \mathcal{H}(a,b,\mathbb{Z})$ we have $\gamma_u = \begin{pmatrix} x_0 + x_1\sqrt{a} & x_2\sqrt{b} + x_3\sqrt{ab} \\ x_2\sqrt{b} - x_3\sqrt{ab} & x_0 - x_1\sqrt{a}\end{pmatrix}$. Letting $x = (x_0, x_1, x_2, x_3)$ and $q_1(x) = q_2(x_0,0,0,x_3) - q_2(0,x_1,x_2,0)$, where $q_2(x) = x_0^2 + ax_1^2 + bx_2^2 + abx_3^2$, we obtain the system

$$\begin{cases} q_1(x) = 1 \\ 2q_2(x) = \|\gamma\|^2 \end{cases} \tag{4}$$

Letting $q_2(x_0,0,0,x_3) = n$ we obtain

$$\begin{cases} x_0^2 + abx_3^2 = n \\ ax_1^2 + bx_2^2 = n-1 \\ \|\gamma\|^2 = 4n-2, n \in \mathbb{N} \end{cases} \tag{5}$$

Taking $a = 2, b = 5$ we have that $n = 1$ or $n \geq 9$. Clearly $\Gamma_i = 1$ and for $n = 9$ we find an element whose bisector Σ separates i and ∞. Using solutions of the system for $n \leq 46$, we obtain a set of generators for a subgroup of finite index. Figure 5 shows the resulting fundamental domain.



Figure 5: Fundamental domain of $S_{2,5}(\mathbb{Z})$.

Theorem 4.11 *The group* $S_{2,5}(\mathbb{Z}) = \langle -1, g(\gamma) \mid g \in \langle \sigma^2, \phi \rangle, \gamma \in X \}$*, where* $X = \{3 + 2i, 2 + 2i + j + k, 3 + 3i + k, 6 + 3j + k\}$*, is a subgroup of finite index of* $\mathrm{SL}_1(\mathcal{H}(2, 5, \mathbb{Z}))$*.*

4.3 Matrix Algebras

In this section we consider groups Γ of finite covolume and at least one ideal vertex. To describe how to implement the DAFC and obtain a fundamental domain we work in $\mathbb{B}^3$ and suppose that j is an ideal vertex. Recall that $S^2 = \partial\mathbb{B}^3$.

Case I: Γ_0 is trivial. First find a $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, with $|a|^2 + |c|^2 = 1$. Since Γ has finite covolume, a finite number of them, those with smallest matrix norm, $\gamma_1, \cdots, \gamma_m$ say, can be chosen such that $V = (S^2 - \{j\}) \bigcap_{1 \le k \le m} \mathrm{Exterior}(\Sigma_{\Psi(\gamma_k)})$ is compact. To know the value of m one just has to know the link of j. Since Γ is of finite covolume, we may use the DAFC to find $\{\gamma_{m+1}.\cdots\gamma_n\}$ such that V is contained in the topological closure of $\bigcup_{1 \le k \le n} \overline{B}_{R_{\Psi(\gamma_k)}}(P_{\Psi(\gamma_k})$. The output, using [12, Theorem II.8.4], is $\langle \gamma_1, \cdots \gamma_n \rangle$, a subgroup of finite index in Γ.

Case II: Γ_0 non-trivial. In this case, proceed as in Case I with $V = (\partial\mathcal{F}_0) \cap (S^2 - \{j\}) \bigcap_{1 \le k \le m} \mathrm{Exterior}(\Sigma_{\Psi(\gamma_k)})$, where $\mathcal{F}_0$ is a fundamental domain of Γ_0. Then $\langle \Gamma_0, \gamma_1, \cdots \gamma_n \rangle$ is of finite index in Γ.

The examples we give here are the Bianchi Groups (see [12, 14]), for which a Ford Fundamental domain is given in [12, Theorem VII.3.4]. As seen in Section 3, the totally geodesic surfaces used in the construction of the Ford domain are exactly the isometric spheres and the planes determined by the generators of Γ_∞. Here we describe a Dirichlet fundamental domain for all d. Later in this section we give a complete proof that the Bianchi groups have finite covolume and describe their ideal vertices. This make this section independent of [12, Chapter VII of] and [40]. As we already saw, for some values of d a Ford domain is also a Dirichlet domain. We shall see one more example of this situation. Note that the Bianchi groups can also be handled as groups commensurable with the unit group of an order in the split quaternion algebra $\mathcal{H}(K)$, $K = \mathbb{Q}(\sqrt{-d})$ and $d \equiv 1, 2 \bmod 4$ or $d \equiv 3 \bmod 4$ and $\mathcal{H}(K)$ not a division ring. All this can be handled as in the previous section (the division assumption in the previous section was only used to guarantee that the groups were cocompact and hence of finite covolume).

Let $\omega = \sqrt{-d}$ if $d \equiv 1, 2 \ mod\ 4$, $\omega = \frac{1+\sqrt{-d}}{2}$ if $d \equiv 3 \ mod\ 4$ and $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma = \mathrm{PSL}(2, \mathbb{Z}[\omega])$. Note that $|a|^2, |b|^2, |c|^2, |d|^2 \in \mathbb{Z}$ and hence Γ is not cocompact. In the ball model $\Gamma_0 = \Gamma \cap \mathrm{SU}(2, \mathbb{C})$, $|a|^2 + |c|^2 = 1$ and thus $a \cdot c = 0$. This implies that a or c is a root of unity, being non-trivial only when $d = 1, 3$. If $d = 1, 3$ then Γ_0 is isomorphic to $C_2 \times C_2, S_3$, respectively and in all other cases $\Gamma_0 = \langle \Psi(\gamma_0) \rangle$, where $\gamma_0 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Since $\gamma_0 \in \Gamma_0$, for all d, we have that a fundamental domain $\mathcal{F}_0$ of Γ is contained in $\{(x, y, z) \in \mathbb{B}^3 \ |z| \geq 0\}$. Since j is an ideal vertex of Γ, we have to find the elements γ such that $j \in \Sigma_{\Psi(\gamma)}$. For these γ´s we have that $|a|^2 + |c|^2 = 1$; if $c(\gamma) \neq 0$ then $a(\gamma_0\gamma) \neq 0$ and $\Sigma_{\Psi(\gamma_0\gamma)} = \Sigma_{\Psi(\gamma)}$. Consequently, we may suppose that $c(\gamma) = 0$ and thus $\gamma \in \Gamma_j$. Denoting by $\mathcal{F}_j$ a fundamental domain of Γ_j acting on S^2, we have that $\mathcal{F} \subset \mathcal{F}_0 \cap \mathcal{F}_j \cap \{(x, y, z) \in \mathbb{B} \ |z \geq 0\}$. So we proved that we may take $\{\gamma_1, \cdots, \gamma_m\} \subset \Gamma_j$ and $V = (\partial \mathcal{F}_j) \cap (\partial \mathcal{F}_0) \cap S^2$.

Using $\eta_0 : \mathbb{H}^3 \to \mathbb{B}^3$ we transfer this information to $\mathbb{H}^3$. In this model $\Gamma_\infty = \{\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \mid ad = 1, a, b, d \in \mathbb{Z}[w]\}$. If $d \notin \{1, 3\}$ then Γ_∞ is isomorphic to a full lattice in $\mathbb{C}$. In any case it is a crystallographic group acting on $\mathbb{C}$. Let $\hat{\mathcal{F}}_\infty$ be a fundamental domain of Γ_∞ acting on $\mathbb{C}$. Then, for all d, $\mathcal{F}_\infty = \{z + rj \in \mathbb{H}^3 \mid z \in \hat{\mathcal{F}}_\infty\}$ is a fundamental domain of Γ_∞ in $\mathbb{H}^3$. Note that $\hat{\mathcal{F}}_\infty$ is the projection on $\partial\mathbb{H}^3$ of $\eta^{-1}(L)$, where L is a link of Γ at $j \in \mathbb{B}^3$.

Lemma 4.12 *1. If* $d \equiv 1, 2 \ mod\ 4$ *then* $\mathcal{F}_\infty = \{z + rj \in \mathbb{H}^3 \mid -\frac{1}{2} \leq Re(z) \leq \frac{1}{2},\ -\frac{\sqrt{d}}{2} \leq Im(z) \leq \frac{\sqrt{d}}{2}\}$

2. *If* $3 < d \equiv 3 \ mod\ 4$ *then* $\mathcal{F}_\infty = \{z + rj \in \mathbb{H}^3 \mid -\frac{1}{2} \leq Re(z) \leq \frac{1}{2},\ -\frac{1+d}{4} \leq Re(z) + \sqrt{d}Im(z) \leq \frac{1+d}{4},\ -\frac{1+d}{4} \leq Re(z) - \sqrt{d}Im(z) \leq \frac{1+d}{4}\}$

3. *If* $d = 3$, *then* $\mathcal{F}_\infty = \{z + rj \in \mathbb{H}^3 \mid 0 \leq Re(z) \leq \frac{1}{2},\ 0 \leq Re(z) + \sqrt{3}Im(z) \leq 1\}$.

4. *If* $d = 1$ *then* $\mathcal{F}_\infty = \{z + rj \in \mathbb{H}^3 \mid -\frac{1}{2} \leq Re(z) \leq \frac{1}{2},\ 0 \leq Im(z) \leq \frac{1}{2}\}$.

Theorem 4.13 *Let* $X = \{\gamma \in \Gamma \mid \Sigma_\gamma \cap \mathcal{F}_\infty \neq \emptyset, \Sigma_\gamma \not\subset \mathbb{B}^3\}$. *Then* $\mathcal{F} = (\mathbb{B}^3)^c \cap \mathcal{F}_\infty \cap \bigcap_{\gamma \in X} Exterior(\Sigma_\gamma)$ *is a Dirichlet fundamental domain for* Γ. *Since* Γ *is of finite covolume,* X *can be chosen to be finite.*

For $d \equiv 1, 2$ mod 4, $\hat{\mathcal{F}}_\infty$ is a rectangle with vertices $\pm\frac{1}{2} \pm \frac{\sqrt{d}}{2}i$ and for $d \equiv 3 \ mod\ 4$ it is a hexagon with vertices $\pm\frac{(d+1)\sqrt{d}}{4d}i$ and $\pm\frac{1}{2} \pm \frac{(d-1)\sqrt{d}}{4d}i$. For $d \neq 3$, all vertices of this hexagon lie on the circle centered at 0 with radius $\frac{(d+1)\sqrt{d}}{4d}$. Hence, for $d \in \{1, 2, 3, 7, 11\}$, $\partial(\hat{\mathcal{F}}_\infty) \subset S_1(0)$ and hence ∞ is the only ideal vertex of Γ.

DAFC was implemented for these groups for $d \equiv 3$ mod 4 (the other cases are treated in the same way.). Let $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, $a = x_0 + y_0 w$, $b = x_1 + y_1 w$, $c = x_2 + y_2 w$, $d = x_3 + y_3 w$, $x = (x_0, x_1, x_2, x_3)$, $y = (y_0, y_1, y_2, y_3)$, $\det(x) = x_0 x_3 - x_1 x_2$ and $J(x) = (x_3, -x_2, -x_1, x_0)$. From this and the condition on $\det(\gamma)$, we obtain the following system of Diophantine equations.

$$\begin{cases} \det(x) - (\frac{d+1}{4})\det(y) = 1 \\ \det(y) + \langle x|J(y)\rangle = 0 \\ \|\gamma\|^2 = \|x\|^2 + (\frac{d+1}{4})\|y\|^2 + \langle x|y\rangle \in \mathbb{N} \end{cases} \tag{6}$$

If $\|\gamma\|^2 = n \in \mathbb{N}$ than $\max\{|a|,|b|,|c|,|d|\} \leq \sqrt{n}$ and hence the equation $\|\gamma\|^2 = n$ has a finite number of solutions (x,y). Consequently, for each $n \in \mathbb{N}$ the system above has a finite number of solutions (x,y) such that $\|\gamma\|^2 = n$.

Theorem 4.14 *For $d \in \{19, 23, 27\}$, the subgroup $\Gamma_{(d)} = \langle \gamma_0, g(X_d) \mid g \in \langle \sigma^2, \tau\rangle\rangle$, where X_d is given below, is of finite index in* $\mathrm{PSL}(2, \mathbb{Z}[w]), w = \frac{1+\sqrt{-d}}{2}$.

$$X_{19} = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -\omega \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1-w & 2 \\ 2 & w \end{pmatrix} \right\},$$

$$X_{23} = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -\omega \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -2+\omega & 3 \\ -1-\omega & -3 \end{pmatrix}, \begin{pmatrix} -3+\omega & 2+\omega \\ -2-\omega & -3+\omega \end{pmatrix} \right\},$$

$$X_{27} = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -\omega \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & -\omega \\ 1-\omega & -3 \end{pmatrix} \right\},$$

The pictures below show a fundamental domain, rotated over ninety degrees, of Γ_d and its projection on $\partial\mathbb{H}^3$ for $d \in \{19, 23, 27\}$, respectively. For $d = 19$ there are no singular points and, by the criteria of Theorem 4.4, the domain is a DF domain (see also [40, Theorem 16.1]). For $d = 23$ note that the ideal vertices are different from the singular points found in a Ford domain (see [40, Lemma 9.10]).



(a) $\Gamma_{(19)}$

(b) $\Gamma_{(19)}$

Figure 6:



(a) $\Gamma_{(23)}$ (b) $\Gamma_{(23)}$

Figure 7:



(a) $\Gamma_{(27)}$ (b) $\Gamma_{(27)}$

Figure 8:

It is worth noting that, for $w = \frac{1+\sqrt{-d}}{2}$, $\gamma = \begin{pmatrix} -\overline{w} & \frac{d-3}{4} \\ 1 & -w \end{pmatrix}$ satisfies that $\Sigma_\gamma = ISO_\gamma$, $P_\gamma = w$

and $R_\gamma = 1$. In general, if $\gamma = \begin{pmatrix} -\overline{w} & b \\ c & -w \end{pmatrix}$, with $b, c \in \mathbb{Z}, bc = \frac{d-3}{4}$, then $\Sigma_\gamma = ISO_\gamma$, $P_\gamma = \frac{w}{c}$ and $R_\gamma = \frac{1}{|c|}$. Similar matrices can be obtained in case $w = \sqrt{-d}$. For small values of d, these matrices, their images by the symmetries and Γ_0, form a set of generators of a subgroup of finite index in the corresponding Bianchi group. The symmetries show that the number of ideal vertices minus one is a multiple of four, in case there are no ideal vertices on the imaginary axis.

Corollary 4.15 *Let* $d \in \{1, 2, 3, 5, 7, 11, 19\}$. *Then the Bianchi group* $PSL(2, \mathbb{Z}[w])$ *has a DF domain.*

Note that the values appearing in the corollary, with the exception of $d = 5$, also appear in others contexts. See for example [4, Proposition 2.3] and [43]. This suggests that having a DF domain is equivalent to other properties. We do not know if these are the only values of d for a DF domain to exist. We now consider the set of cusps, C_Γ, of a Bianchi group Γ. For this we shall first reprove that Γ is of finite covolume and hence geometrically finite. In what follows $K = \mathbb{Q}(\sqrt{-d})$, $\mathfrak{o}_K$ is the ring of integers of K and $\mathrm{Pic}(K)$ is the Picard group of K.

Proposition 4.16 *Let* Γ *be a Biachi group and* $\mathcal{F}$ *a Poincaré fundamental domain of* Γ. *Then the following hold.*

1. Γ *is of the first kind.*
2. *The ideal points of* $\mathcal{F}$ *are contained in* $\mathbb{P}^1(K)$.
3. *If* $z = \frac{\alpha}{\beta} \in K, \alpha, \beta \in \mathfrak{o}_K$ *is an ideal point, then* $\langle \alpha, \beta \rangle$ *is a proper ideal of* $\mathfrak{o}_K$.

Proof. The element $\gamma \in \Gamma$, with $a(\gamma) = c(\gamma) = d(\gamma) = 1, b(\gamma) = 0$, is parabolic and fixes 0. Let $\alpha, \beta \in \mathfrak{o}_K$ be such that the ideal $\langle \alpha, \beta \rangle = \mathfrak{o}_K$. Consider $\gamma_1 \in \Gamma$ such that $b(\gamma_1) = \alpha$ and $d(\gamma_1) = \beta$. Then $\frac{\alpha}{\beta}$ is the fixed point of $\gamma_1 \gamma \gamma_1^{-1}$ and hence, by [12, Proposition VII.2.7], the set of fixed points of parabolic elements of Γ is dense in $\mathbb{P}^1(\mathbb{C})$. Applying [33, Theorem XII.1.5] we get that $L(\Gamma) = \mathbb{P}^1(\mathbb{C})$ and so Γ is of the first kind.

Let $z \in \mathbb{C}$, $z \notin K$. By [12, Proposition VII.2.7], there exists a sequence of distinct elements $z_n = \frac{\alpha_n}{\beta_n} \in K$ such that $|z - z_n| < |\beta_n|^{-1}$, $\langle \alpha_n, \beta_n \rangle = \mathfrak{o}_K$. Consider $\gamma_{+n} = \begin{pmatrix} 1 + \alpha_n \beta_n & -\alpha_n^2 \\ \beta_n^2 & 1 - \alpha_n \beta_n \end{pmatrix} \in \Gamma$. We have that, for sufficiently large n,

$$|P_{\gamma_{+n}} - \hat{P}_{\gamma_{+n}}| = \frac{2|\mathrm{Re}(\alpha_n \beta_n)|}{|\beta_n|^2(|1 + \alpha_n \beta_n|^2 + |\beta_n|^4 - 1)} \leq \frac{2||\alpha_n \beta_n|}{|\beta_n|^2(|1 + \alpha_n \beta_n|^2 + |\beta_n|^4 - 1)} \leq |\beta_n|^{-2},$$

$|\hat{P}_{\gamma_{+n}} - z_n| = |\beta_n|^{-2}$ and, since by Lemma 3.8, $\hat{P}_{\gamma_{+n}} \in \mathrm{Interior}(\Sigma_{\gamma_{+n}})$, $|P_{\gamma_{+n}} - z| \leq 2|\beta_n|^{-2}$. We will estimate $R^2_{\gamma_{+n}}$ from which it will follow that, for sufficiently large n, $z \in \mathrm{Interior}(\Sigma_{\gamma_{+n}})$. Indeed, for sufficiently large n and remembering that $|\lambda|^2 \in \mathbb{N}$ for $\lambda \in \wr_K$, we have, by Proposition 3.3, that

$$R^2_{\gamma_{+n}} = \frac{(|1 + \alpha_n \beta_n|^2 + |\beta_n|^4 - 1)^2 + (|1 + \alpha_n \beta_n|^2 + |\beta_n|^4)(|\alpha_n|^4 + |1 - \alpha_n \beta_n|^2) - 1}{(|1 + \alpha_n \beta_n|^2 + |\beta_n|^4)(|1 + \alpha_n \beta_n|^2 + |\beta_n|^4 - 1)} \geq$$

$$\frac{|\alpha_n|^4 + |1 - \alpha_n \beta_n|^2}{|1 + \alpha_n \beta_n|^2 + |\beta_n|^4 - 1} = |z_n|^4 \cdot \frac{1 + |\alpha_n^{-2} + z_n^{-1}|^2}{1 + |\beta_n^{-2} + z_n|^2 - |\beta_n|^{-4}}.$$

Since (z_n) converges to z, we have that $\liminf R_{\gamma_{+n}} > 0$, i.e., (R_{γ_n}) is bounded away from 0. Consequently, for sufficiently large n, $z \in \mathrm{Interior}(\Sigma_{\gamma_{+n}})$ and this proves the second item.

Suppose that $\langle \alpha, \beta \rangle = \mathfrak{o}_K$ and set $\gamma = \begin{pmatrix} x & -y \\ -\beta & \alpha \end{pmatrix} \in \Gamma$. Then, by Corollary 3.7 and Lemma 3.8, $z = \hat{P}_\gamma \in \mathrm{Interior}(\Sigma_\gamma)$. ▮

Let $p = \frac{\alpha}{\beta} \in K$ corresponds to a non-trivial element of $\mathrm{Pic}(K)$ and, following [12, Definition VII.3.5], define $\gamma_+ = \begin{pmatrix} 1+\alpha\beta & -\alpha^2 \\ \beta^2 & 1-\alpha\beta \end{pmatrix}$, $\gamma_- = \begin{pmatrix} 1+w\alpha\beta & -w\alpha^2 \\ w\beta^2 & 1-w\alpha\beta \end{pmatrix}$ and $\gamma = \begin{pmatrix} \alpha\beta & 1 \\ \beta^2 & 0 \end{pmatrix}$. Then $\langle \gamma_+, \gamma_- \rangle < \Gamma_p$, $\gamma^{-1}\gamma_+\gamma = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, $\gamma^{-1}\gamma_-\gamma = \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix}$ and hence $\langle \gamma_+, \gamma_- \rangle$ is isomorphic to $\mathbb{Z}^2$. The intersection of the union of the interiors of the bisectors $\Sigma_{\gamma_+^{\pm 1}}$ and $\Sigma_{\gamma_-^{\pm 1}}$ with $\partial(\mathbb{H}^3)$, contains a punctured disk of $\partial\mathbb{H}^3$ centered in p. This is easily seen using Lemma 3.12. From this it follows easily that a cusp region has finite covolume. We now prove that all ideal vertices of Γ, except for ∞, arise in this way.

Theorem 4.17 *Let Γ be a Bianchi group. Then the following hold.*

1. *Γ has finite covolume and is geometrically finite.*
2. *If $z \in \mathbb{P}^1(\mathbb{C})$ is an ideal vertex of Γ, it corresponds to a non-trivial element of* $\mathrm{Pic}(K)$ *and* $1 \leq |z|$.
3. *A fundamental domain $\mathcal{F}$ of Γ has $|\mathrm{Pic}(K)|$ ideal vertices.*

Proof. By Proposition 4.16 and [12, Theorem VII.2.4] it follows that $\mathcal{F}$ has $|\mathrm{Pic}(K)|$ ideal points. By the observation before the theorem, each cusp region has finite covolume from which it easily follows that $vol(\Gamma)$ is finite. For every ideal vertex $z_i, 1 \leq i \leq |\mathrm{Pic}(K)|$, let V_{z_i} be the interior of the projection on $\partial(\mathbb{H}^3)$ of a horoball centered at z_i. Let $V = \hat{\mathcal{F}}_\infty - \bigcup_i V_{z_i}$. Then V is compact and Proposition 4.16 shows that it can be covered by a finite number of bisectors. Hence Γ is geometrically finite. (This also follows by a result of Garlan and Raghunathan, [12, Theorem II.2.7].)

By Proposition 4.16 an ideal vertex is of the form $z = \frac{\alpha}{n}$ with $n \in \mathbb{N}, \alpha \in \mathbb{Z}[w]$ and with $I = \langle n, \alpha \rangle$ a proper non-zero ideal of $\mathbb{Z}[w]$. Suppose it were principal, i.e., $\langle n, \alpha \rangle = I = \langle \beta \rangle$. We may write $\beta = nx + y\alpha$, $n = u\beta$ and $\alpha = v\beta$, with $x, y, u, v \in \mathbb{Z}[w]$. From this it follows that $xu + yv = 1$ and hence $z = \frac{v}{u}$ with $\langle u, v \rangle = \mathbb{Z}[w]$, contradicting Proposition 4.16. ▮

The proofs of Proposition 4.16 and Theorem 4.17 suggests that a Bianchi group is virtually generated by parabolic elements. In case $d = 3$ a subgroup of finite index generated by parabolic elements is the figure-eight knot group.

We give more information about an ideal vertex $z \in \mathbb{C}$ of the previous corollary which will allow the determination of all the ideal vertices. Write $t = \gcd(n, |\alpha|^2 - 1)$, $n = ts$ and $|\alpha|^2 - 1 = tm$. If we set $\gamma_1 = \begin{pmatrix} \overline{\alpha} & -m \\ -t & \alpha \end{pmatrix}$, then $sz = P_{\gamma_1}$ and $\Sigma_{\gamma_1} = \mathrm{ISO}_{\gamma_1}$. Hence, since z corresponds to an ideal vertex of the fundamental domain $\mathcal{F}$ of Γ (given in Theorem 4.13), we have that $s \neq 1$, i.e., n is not a divisor of $|\alpha|^2 - 1$. By [9, Theorem IV.4.24], I can be represented as $a\mathbb{Z} + (b + cw)\mathbb{Z}$, where

$a, b, c \in \mathbb{Z}$, where $a > 0, c > 0, 0 \le b < a$, $c \mid b$, $c \mid a$ and $ac \mid |b + cw|^2$. Hence we may write $b = cr$ and $a = cs$. From this we conclude that $z = \frac{\alpha}{n} = \frac{b+cw}{a} = \frac{r+w}{s}$, with $r, s \in \mathbb{Z}$, $s > 0$, $0 \le r < s$ and $s \mid |r + w|^2$. In particular, z corresponds to the element $[s, r + w] \in \mathbb{P}^1(\mathbb{C})$. Since $z \in \hat{\mathcal{F}}_\infty$, it follows that $|\mathrm{Re}(r + w)| \le \frac{s}{2}$, $s \le |r + w|$ and $\frac{3s^2}{4} \le |\mathrm{Im}(r + w)|^2 \le \frac{ds^2}{4}$. By [12, Theorem VII.2.4], Γ has $h = |\mathrm{Pic}(K)|$ non Γ-equivalent ideal vertices. Hence we have described all the ideal vertices of $\mathcal{F}$ and thus we have proved the following theorem which is due to Swan (see [40, Corollary 7.6]) in the context of a Ford domain. In case $d \in \{1, 2, 3, 7, 11, 19\}$, then the only ideal vertex of Γ is ∞.

Theorem 4.18 *The ideal vertices of the Bianchi group* $\Gamma = \mathrm{PSL}(2, \mathbb{Z}[w])$ *are* ∞ *and the points* $g(z)$, $z = \frac{r+w}{s} \in \hat{\mathcal{F}}_\infty$, *with* $g \in \langle \sigma^2, \tau \rangle$, $r, s \in \mathbb{Z}$, $s > 0$, $0 \le r < s$, $0 \le 2|\mathrm{Re}(r + w)| \le s$, $s \mid |r + w|^2$, $s \le |r + w|$ *and* $|\mathrm{Im}(r + w)|^2 \le \frac{ds^2}{4}$. *More precisely we have:*

1. *For* $d \equiv 1, 2 \bmod 4$, *we have that* $4 \le s^2 \le \frac{4d}{3}$ *and* $0 \le r \le \frac{s}{2}$.
2. *For* $d \equiv 3 \bmod 4$, *we have that* $1 < s \le \frac{2+\sqrt{4+3d}}{3}$ *and* $0 \le r \le \frac{s-1}{2}$.

It is well known that the figure-eight knot group $\Gamma_8 = \langle \gamma_1, \gamma_2 \rangle$ is generated by $\gamma_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $\gamma_2 = \begin{pmatrix} 1 & 0 \\ w & 1 \end{pmatrix}$, $w = \frac{-1+\sqrt{3}}{2}$ (see [12, 41]). A matrix presentation for Γ_8 was first described in [36] where also a fundamental domain was given (see also [13, 31]). Looking at the generating set, and having in mind the criterium of Theorem 4.4, one might think that Γ_8 has a DF domain. However we show that this is not the case as it is also not the case for the group defining the hyperbolic structure on the complement of the Whitehead Link. To see this we will use the results of Section 3 and [33, Figures 10.3.3 and 10.3.14] which show the gluing patterns for the figure-eight knot complement and the Whitehead Link complement. We stick to the notation used in the figures of [33]. In the first case face B is a vertical plane which is mapped to B' an Euclidean sphere and face D is an Euclidean sphere which is mapped to a vertical plane. In the case of the Whitehead Link face A is a vertical plane which is mapped to A' an Euclidean sphere. In fact, in this case all vertical planes are mapped to Euclidean spheres and vice versa. This violates a basic property of a DF domain: *If* $\mathcal{F}$ *is a fundamental domain for* Γ *and* γ *is a side-pairing transformation such that* $\infty \in \Sigma_\gamma$ *then* $\gamma \in \Gamma_\infty$. This follows from the proof of Theorem 4.4. We now describe a Poincaré fundamental domain $\mathcal{F}$ for Γ_8. We shall make use of the matrix presentation given in [36]. We could use the Cayley graph and apply DAFC but, in this case, it is very easy to get a set of generators needed to construct a Poincaré fundamental domain.

Indeed, if we set $\gamma_3 = \begin{pmatrix} 0 & w \\ -w^2 & 1 - w \end{pmatrix}$, $\gamma_4 = \begin{pmatrix} 0 & -w \\ w^2 & 1 - w \end{pmatrix}$ and $\gamma_5 = \begin{pmatrix} 1 & -1 \\ -w & 1 + w \end{pmatrix}$ then, using the results of Section 3, one can obtain, by hand, the bisectors of these elements and of their inverses. Note that Σ_{γ_3} is a vertical plane but $\Sigma_{\gamma_3{-1}}$ is an Euclidean sphere. This is also the case for γ_4. As explained, this can be deduced from the gluing patterns and the results of Section 3. The pictures below show the projection of the link and the fundamental domain.



(a) Link and bisectors (b) Poincaré Fundamental domain

Figure 9:

We now use the results of Section 3 to obtain a matrix representation of the group Γ defining the hyperbolic structure on the Whitehead Link complement. We will use the regular octahedron T of [33, Figure 10.3.12], the gluing pattern [33, Figure 10.3.13] and the side-pairing transformations explained on [33, Page 454]. The notation of [33, Page 454] is maintained. We will now work in both $\mathbb{B}^3$ and $\mathbb{H}^3$ and use freely the results of Section 3. Set $\Psi(\gamma_1) = g_A$, $\Psi(\gamma_2) = g_B$, $\Psi(\gamma_3) = g_C$ and $\Psi(\gamma_4) = g_D$. Clearly $\Gamma = \langle \gamma_1, \gamma_2, \gamma_3, \gamma_4 \rangle$. We have that $\Sigma_{\Psi(\gamma_1)} = A$, $\Sigma_{\Psi(\gamma_2)} = B$, $\Sigma_{\Psi(\gamma_3)} = C$ and $\Sigma_{\Psi(\gamma_4)} = D$. Note that η_0^{-1} fixes pointwise the unit circle $\mathbb{B}^3 \cap \partial(\mathbb{H}^3)$ and $\mathcal{F} = \eta_0^{-1}(T)$ is a Poincaré fundamental domain in $\mathbb{H}^3$ of Γ. We can now easily describe explicitly all the γ_k's. Since they are Möbius transformations we only have to give them at three points. We use ordered triples to write them. Using the gluing pattern we easily deduce that $\gamma_1 : (i, 1, 0) \mapsto (-i, \infty, 1)$, $\gamma_2 : (i, -1, 0) \mapsto (i, \infty, 1)$, $\gamma_3 : (-1, -i, 0) \mapsto (\infty, i, -1)$ and $\gamma_4 : (-i, 1, 0) \mapsto (-i, \infty, -1)$. Using cross-ratio, we find that $\gamma_1 = \lambda \begin{pmatrix} -2-i & i \\ -i & i \end{pmatrix}$, $\gamma_2 = \lambda \begin{pmatrix} 2-i & -i \\ -i & -i \end{pmatrix}$, $\gamma_3 = \lambda \begin{pmatrix} -2-i & -i \\ i & i \end{pmatrix}$ and $\gamma_4 = \lambda \begin{pmatrix} 2-i & i \\ i & -i \end{pmatrix}$, where $\lambda = \frac{\sqrt{2i}}{2}$. Note that $\gamma_3 = \sigma^2(\gamma_1)$ is elliptic, $\gamma_4 = \sigma^2(\gamma_2)$ is parabolic and $\phi(\mathcal{F}) = \mathcal{F}$. One can check, using the formulas of Section 3, that the γ_k's are indeed the side-pairing transformations of $\mathcal{F}$. Clearly $\langle \gamma_1, \gamma_2, \sigma^2 \rangle$ is a Kleinian group containing Γ as a subgroup of index two, i.e., the Whitehead link complement is the double cover of an orbifold. This is also known for the figure-eight knot complement group as $\langle \Gamma_8, \delta \rangle = \mathrm{PSL}(2, \mathbb{Z}[w])$, with $d = 3$.

From the gluing pattern given in [33, Figure 10.3.22], it is easy to deduce that the group giving the hyperbolic structure of the Borrromean rings complement has a fundamental domain in which the basic property of a DF domain is satisfied. Using the presentation given in [37] and the results of

Section 3, one can construct a Poincaré fundamental domain of the Borrromean rings complement group.

5 Group Rings

The last chapter of [38] contains a list of open problems, among which Problem 17 and 23, asking for generators and relations or generators of a subgroup of finite index of $\mathcal{U}(\mathbb{Z}G)$ for finite groups G.

The former has been done for very few groups, basically for all groups of order at most 16 (see for example [38]).

Bass and Milnor introduced the Bass cyclic units and proved that they generate a subgroup of finite index in $\mathcal{U}(\mathbb{Z}G)$ for G a finite abelian group ([5]). Ritter and Sehgal introduced the bicyclic units and proved that, for many finite groups G, the Bass cyclic and bicyclic units generate a subgroup of finite index in $\mathcal{U}(\mathbb{Z}G)$ (see [38]).

A simple algebra A is said to be of *exceptional type* if it is either a non-commutative division algebra (which is not a totally definite quaternion algebra), or $M(2,\mathbb{Q})$, or $M(2,\mathbb{Q}(\sqrt{-d})), d > 0$, or $M(2,D)$, where D is a rational quaternion division algebra $\mathcal{H}(a,b,\mathbb{Q})$. Let G be a finite group with no exceptional Wedderburn components. For every primitive central idempotent e_i of $\mathbb{Q}G$ for which $(\mathbb{Q}G)e_i$ is not a division ring, let f_i be a non-central idempotent in $(\mathbb{Q}G)e_i$ and n_{f_i} a positive integer so that $n_{f_i}f_i \in \mathbb{Z}G$. Then, in [17], Jespers and Leal gave the following extention of the results of Ritter and Sehgal described above (the proof ultimately relies on the congruence theorem of Bass-Milnor-Serre [6], Vasserstein [42] and Bak-Rehmann [3]). The group generated by the Bass cyclic units and the units of the form $1 + n_{f_i}^2 f_i h(1-f_i)$ and $1 + n_{f_i}^2(1-f_i)hf_i$, $h \in G$, is of finite index in $\mathcal{U}(\mathbb{Z}G)$. If, moreover, Ge_i is not fixed point free then one may take $f_i = \frac{1}{o(g_i)}\hat{g}_i$, for some $g_i \in G$. Hence, if G has no non-abelian fixed point free images then the Bass cyclic and the bicyclic units generate a subgroup of finite index in $\mathcal{U}(\mathbb{Z}G)$.

The numbers n_{f_i} could be replaced by any number so that $n_{f_i}^2 f_i \mathbb{Z}G(1-f_i) \subseteq \mathbb{Z}G$. In the proof of this result it is shown that the group generated by the units of the form $1 + n_{f_i}^2 f_i h(1-f_i)$ and $1 + n_{f_i}^2(1-f_i)hf_i$ contain a subgroup of finite index in $\mathrm{SL}(n_i, Ø_i)$ provided $\mathbb{Q}Ge_i$ is not of exceptional type and $n_i \geq 2$. Although they are not central, the Bass cyclic units are needed to obtain a subgroup of finite index in the center.

In [20] the rational representations of a finite nilpotent group G have been explicitly described by exhibiting a set of matrix units of each simple component of $\mathbb{Q}G$. In particular, a precise description is given of when a simple component $\mathbb{Q}Ge$ (e a primitive central idempotent) is of exceptional type (this also follows from [18]): *These $\mathbb{Q}Ge$'s are one of the following algebras: $\mathcal{H}(\mathbb{Q}(\xi_{2^{n-1}} + \xi_{2^{n-1}}^{-1}))$, $\mathcal{H}(\mathbb{Q}(\xi_n))$, $M(2,\mathbb{Q})$, $M(2,\mathbb{Q}(\sqrt{-2}))$, $M(2,\mathbb{Q}(i))$, $M(2,\mathbb{Q}(\sqrt{-3}))$ or $M(2,\mathcal{H}(\mathbb{Q}))$. Let G_e denote the stabilizer of e in G.. For each possible form of $\mathbb{Q}Ge$, a description of G/G_e is given and, in case $\mathbb{Q}Ge$ is a matrix algebra, a complete set of matrix units is also given* (see also [18]).

As a consequence of these results, still supposing G nilpotent, it follows that if G/G_e is a non-commutative fixed point free group and $\mathbb{Q}Ge$ is a non-commutative division algebra then $G/G_e = Q_8 \times C_n = \langle a,b,c \mid a^4 = 1,\ a^2 = b^2,\ ba = a^{-1}b,\ ca = ac, cb = bc\rangle$ with n so that 2 has odd order in $\mathcal{U}(\mathbb{Z}_n)$. Let G be a finite group. For a subgroup H of G we put $\hat{H} = \sum_{h\in H} h \in \mathbb{Z}G$ and $\widetilde{H} = \frac{\hat{H}}{|H|}$.

Proposition 5.1 *Let G be a finite nilpotent group and e a primitive central idempotent of $\mathbb{Q}G$ so that G/G_e is a non-commutative fixed point free group and $\mathbb{Q}Ge$ is not a division algebra. Then*

$\mathbb{Q}Ge = M(2, \mathbb{Q}(\xi_n))$ and $e = \widetilde{G_e}\frac{1}{2}(1-a^2)\left(1-\frac{1}{n}(\hat{c})\right)$. Furthermore, $\mathbb{Q}Ge$ has the following set of matrix units:

$$E_{11} = e\tfrac{1}{2}(1+xa+yab), \qquad E_{12} = e\tfrac{1}{2}(1+xa+yab)b,$$

$$E_{21} = eb^{-1}\tfrac{1}{2}(1+xa+yab), \quad E_{22} = e\tfrac{1}{2}(1-xa-yab).$$

where $x = \frac{1}{2}(\alpha + b\alpha b^{-1})$, $y = b^3\frac{1}{2}(\alpha - b\alpha b^{-1}) \in \mathbb{Z}\langle c^{n/p}\rangle$, and $\alpha = \prod_{k=0}^{m-1}\left(1 + bc^{(n/p)2^k}\right)$ and p is a prime divisor of n so that $2^p \equiv -1$ (mod p).

Proof. It follows from [15, Lemma, page 153] that the elements x and y satisfy $x^2+y^2+1=0$ in $\mathbb{Q}\langle c^{n/p}\rangle e \cong \mathbb{Q}(\xi_p)$. Hence, the result follows from [20, Theorem 4.5]. ▮

Let G be a finite group and assume N is a normal subgroup of G and e is a primitive central idempotent of $\mathbb{Q}G\hat{N}$ so that $\mathbb{Q}G\hat{N}e = M(2,F)$ with F a field. Suppose $\O$ is an order in F and $E_{11}, E_{12}, E_{21}, E_{22}$ is a set of matrix units of $M(2,F)$ belonging to $\mathbb{Q}G\hat{N}e$, then, for a positive integer m, we denote by $\Gamma_{2,N}(m\O)$ the congruence group of level m, that is

$$\Gamma_{2,N}(m\O) = \left(1 + \sum_{1\le i,j\le 2}(m\O)E_{ij}\right) \cap \mathrm{SL}(2,\O) = (1 + M(2, m\O)) \cap \mathrm{SL}(2,\O).$$

There exists a positive integer m so that $m\mathcal{O}E_{ij} \subseteq \mathbb{Z}G$, for all i,j, and thus for such an m we have that $\Gamma_{2,N}(m\mathcal{O})$ is a subgroup of $\mathcal{U}(\mathbb{Z}G)$.

In [19], it was shown that if G is a nilpotent finite group of nilpotency class n then the group $\langle b_{(n)} \mid b \text{ a Bass cyclic unit}\rangle$ is of finite index in $Z(\mathcal{U}(\mathbb{Z}G))$. Here $b_{(n)}$ is defined recursively as follows. We denote by Z_i the i-th centre of G. For any $x \in G$ and Bass cyclic unit $b \in \mathbb{Z}\langle x\rangle$ put $b_{(1)} = b$, and, for $2 \le i \le n$, put $b_{(i)} = \prod_{g\in Z_i} b_{(i-1)}^g$, where $\alpha^g = g^{-1}\alpha g$ for $\alpha \in \mathbb{Z}G$.

Theorem 5.2 *Let G be a nilpotent finite group of nilpotency class n. Assume that the rational group algebra $\mathbb{Q}G$ does not have simple components that are non-commutative division algebras (except totally definite quaternion algebras) or $M(2, \mathcal{H}(a,b,\mathbb{Q}))$. Then the group generated by the following units is of finite index in $\mathcal{U}(\mathbb{Z}G)$:*

1. *$b_{(n)}$, with b a Bass cyclic unit in $\mathbb{Z}G$,*
2. *the bicylic units in $\mathbb{Z}G$,*
3. *the groups $\Gamma_{2,N}(m\O)$ (where for the respective congruence groups we use the matrix units described in [20]) with N a normal subgroup of G so that*
   - *(a) $\O = \mathbb{Z}$, $m = 8|N|$ and $G/N = D_8$,*
   - *(b) $\O = \mathbb{Z}[\sqrt{-2}]$, $m = 8|N|$ and $G/N = D_{16}^-$,*
   - *(c) $\O = \mathbb{Z}[i]$, $m = 8|N|$ and $G/N = D_{16}^+$,*
   - *(d) $\O = \mathbb{Z}[\sqrt{-3}]$, $m = 24|N|$ and $G/N = D_8 \times C_3$,*
   - *(e) $\O = \mathbb{Z}[\sqrt{-3}]$, $m = 24|N|$ and $G/N = Q_8 \times C_3$,*
   - *(f) $\O = \mathbb{Z}[i]$, $m = 32|N|$ and $G/N = D^+$,*

4. *$u_{g,N} = 1 + E_{11}gE_{22}$ and $u'_{g,N} = 1 + E_{22}gE_{11}$, when $g \in G$ and N a normal subgroup of G so that $G/N = Q_8 \times C_n$ such that 2 has even order in $\mathcal{U}(\mathbb{Z}_n)$ (with E_{11} and E_{22} as stated in Proposition 5.1 and $N = G_e$).*

Proof. As mentioned earlier, the units listed in (1) generate a subgroup of finite index in $\mathcal{Z}(\mathcal{U}(\mathbb{Z}G))$. Note that $\mathcal{H}(\mathbb{Q}(\xi_n + \xi_n^{-1}))$ is a totally definite quaternion algebra and hence (see [38, Lemma 21.3]) the unit group of an order $\emptyset$ in this algebra is central-by-finite. Hence the group $\langle b_{(n)} \mid b$ a Bass cylcic unit$\rangle$ contains a subgroup of finite index in $\emptyset$. Hence, by the previously mentioned results of [17], we need to show that the group generated by the proposed units contains a subgroup of finite index in $\mathrm{SL}(n, \emptyset)$ for each simple component $\mathbb{Q}Ge = M(n, D)$ (e a primitive central idempotent of $\mathbb{Q}G$ and D a division algebra) so that $\mathbb{Q}Ge$ is not commutative. Again, by the results of [17] mentioned above, the group generated by units listed in (2) contains such a subgroup provided $\mathbb{Q}Ge$ is not of exceptional type and Ge is not fixed point free. If $\mathbb{Q}Ge$ is a division algebra, and thus a totally definite quaternion algebra by the assumptions, then we already know this by the first part of the proof. If $G/G_e \cong Ge$ is a fixed point free group then, because of Proposition 5.1, the units listed in (4) provide us with the required subgroup. The units listed in (3) take care of all other components. ∎

In [15] different units were used for (4) and in [18] a more complicated construction was given of units that played the role of those listed in (3). In general, the result fails if one does not include the units listed in (3) (we refer the reader to [38, Section 25]).

Generators for the congruence groups listed in Theorem 5.2.3 can be computed using DAFC. In case the simple component is $M(2, \mathbb{Q})$ one can also use Farey symbols (see [11])

The groups listed in Theorem 5.2.(3) (with for example $|N| = 1$) are of finite covolume (coarea) but not cocompact (because of Corollary 3.7) and a set of generators can be calculated using the DAFC. However, it gives a too large set of generators to be listed here. As a matter of example, we therefore restrict ourselves to determine a set of generators for a subgroup of finite index for each of the following groups: $\Gamma_2(8\mathbb{Z})$, $\Gamma_2(2\mathbb{Z}[\sqrt{-2}])$ and $\Gamma_2(2\mathbb{Z}[i])$.

Let $n_0 \in \mathbb{N}$. In $\mathrm{PSL}(2, \mathbb{Z})$ consider the discrete Fuchsian subgroup $\Gamma_2(n_0\mathbb{Z})= \{\gamma = \begin{pmatrix} 1 + n_0 a & n_0 b \\ n_0 c & 1 + n_0 d \end{pmatrix}, a, b, c, d \in \mathbb{Z}\}$. Write $\gamma = 1 + n_0\hat{\gamma}$, $\hat{\gamma} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then $\|\gamma\|^2 = 2 + n_0^2(\|\hat{\gamma}\|^2 + \frac{2}{n_0}\mathrm{tr}(\hat{\gamma}))$ and $\det(\gamma) = 1$ if and only if $\mathrm{tr}(\hat{\gamma}) + n_0 \det(\hat{\gamma}) = 0$. Hence, it follows that $\|\gamma\|^2 = 2 + n_0^2(\|\hat{\gamma}\|^2 - 2\det(\hat{\gamma}))$.

So we obtain the following defining system of algebraic equations for $\Gamma_2(n_0\mathbb{Z})$.

$$\begin{cases} \|\hat{\gamma}\|^2 - 2\det(\hat{\gamma}) = n \\ \mathrm{tr}(\hat{\gamma}) + n_0 \det(\hat{\gamma}) = 0 \\ \|\gamma\|^2 = 2 + n_0^2 n, n \in \mathbb{N} \end{cases} \tag{7}$$

Next, working in $\mathbb{B}^2$, we look for elements γ whose isometric circle passes through i. In this case, these are exactly the elements stabilizing i. Thus, for such elements γ, we have that $(1 + n_0 a)^2 + n_0^2 c^2 = 1$ and therefore $\gamma = \begin{pmatrix} 1 & n_0 b \\ 0 & 1 \end{pmatrix}$. Hence, their defining bisectors, in $\mathbb{H}^2$, which are vertical lines, are given by the equation $Re(n_0 bz) + \frac{n_0^2 b^2}{2} = 0$ or equivalently $x + \frac{n_0 b}{2} = 0$. The bisector corresponding to such an element of smallest norm is the line $x + \frac{n_0}{2} = 0$. We use the DAFC to cover the compact region $[\frac{-n_0}{2} \leq x \leq \frac{n_0}{2}]$, up to a finite number of points, which are precisely the remaining ideal vertices of Γ. If $n_0 = 2$, we get the group whose orbifold is the trice punctured sphere mentioned in the beginning of the previous section. For $n_0 = 8$, we get the compact region $[-4, 4]$ which is covered using solutions of the above system with $n \leq 2304$. This is depicted in Figure 10.

Theorem 5.3 $\tilde{\Gamma}_2(8\mathbb{Z}) = \langle g(\gamma) \mid g \in \langle \sigma^2, \phi \rangle, \gamma \in X \rangle < \Gamma_2(8\mathbb{Z})$, *where* X *is given by the table below, is of finite index.*

a	*b*	*c*	*d*	*a*	*b*	*c*	*d*	*a*	*b*	*c*	*d*
0	*0*	*1*	*0*	*-3*	*1*	*-9*	*3*	*8*	*-6*	*-11*	*8*
-1	*1*	*-1*	*1*	*5*	*-2*	*-8*	*3*	*13*	*-8*	*8*	*-5*
2	*-1*	*4*	*-2*	*-10*	*7*	*-3*	*2*	*8*	*-3*	*-22*	*8*
-7	*2*	*3*	*-1*	*6*	*-4*	*9*	*-6*	*16*	*-26*	*-10*	*16*
-4	*3*	*5*	*-4*	*11*	*-4*	*8*	*-3*	*21*	*-8*	*34*	*-13*
7	*-2*	*-4*	*1*	*-4*	*1*	*15*	*-4*	*-16*	*6*	*42*	*-16*
-5	*2*	*7*	*-3*								



Figure 10: Fundamental domain of $\tilde{\Gamma}_2(8\mathbb{Z})$.

The defining system of Diophantine equations to describe the groups (and to obtain generators for these groups) in the next theorem are exactly the same as before, i.e.,

$$\begin{cases} \|\hat{\gamma}\|^2 - 2\det(\hat{\gamma}) = n \\ \operatorname{tr}(\hat{\gamma}) + n_0 \det(\hat{\gamma}) = 0 \\ \|\gamma\|^2 = 2 + n_0^2 n, n \in \mathbb{N} \end{cases} \tag{8}$$

only the entries of $\hat{\gamma}$ are now in $\mathbb{Z}[\sqrt{-d}]$. This is the case for any congruence subgroup which we still have to deal with.

Theorem 5.4 *1. The subgroup* $\tilde{\Gamma}_2(2\mathbb{Z}[\sqrt{-2}]) = \langle -1, g(X_1) \mid g \in \langle \sigma^2 \circ \tau, \phi \rangle \rangle < \Gamma_2(2\mathbb{Z}[\sqrt{-2}])$, *where* X_1 *is given below, is of finite index .*

2. The subgroup $\tilde{\Gamma}_2(2\mathbb{Z}[i]) = \langle -1, g(X_2) \mid g \in \langle \sigma^2 \circ \tau, \phi \rangle \rangle < \Gamma_2(2\mathbb{Z}[i])$,*where* X_2 *is given below, is of finite index . Its fundamental domain is a DF domain.*

$$\begin{aligned} X_1 \;=\; & \left\{ \begin{pmatrix} 1 & -2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -2\sqrt{-2} \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ -2\sqrt{-2} & -1 \end{pmatrix}, \begin{pmatrix} -3 & 2\sqrt{-2} \\ -2\sqrt{-2} & -3 \end{pmatrix}, \right. \\ & \begin{pmatrix} 1-2\sqrt{-2} & 2\sqrt{-2} \\ -2\sqrt{-2} & -1-\sqrt{-2} \end{pmatrix}, \begin{pmatrix} -1+2\sqrt{-2} & 2 \\ 4 & -1-\sqrt{-2} \end{pmatrix}, \begin{pmatrix} -1+2\sqrt{-2} & 4 \\ 2 & -1-\sqrt{-2} \end{pmatrix}, \\ & \begin{pmatrix} -3+2\sqrt{-2} & 4 \\ 4 & -3-2\sqrt{-2} \end{pmatrix}, \begin{pmatrix} -3-2\sqrt{-2} & 2\sqrt{-2} \\ -4\sqrt{-2} & -3+2\sqrt{-2} \end{pmatrix}, \\ & \left. \begin{pmatrix} -3-2\sqrt{-2} & 4\sqrt{-2} \\ -2\sqrt{-2} & -3+2\sqrt{-2} \end{pmatrix}, \begin{pmatrix} 5-2\sqrt{-2} & 4\sqrt{-2} \\ -4\sqrt{-2} & 5+2\sqrt{-2} \end{pmatrix} \right\}, \end{aligned}$$

$$X_2 = \left\{ \begin{pmatrix} 1 & -2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -2i \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ -2i & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix}, \begin{pmatrix} -1+2i & 2 \\ 2 & -1-2i \end{pmatrix}, \begin{pmatrix} -1-2i & 2i \\ -2i & -1+2i \end{pmatrix} \right\}.$$

A fundamental domain and its projection on $\partial\mathbb{H}$ is given below for the two groups above.



(a) $\tilde{\Gamma}_2(2\mathbb{Z}[\sqrt{-2}])$ (b) $\tilde{\Gamma}_2(2\mathbb{Z}[\sqrt{-2}])$

Figure 11:



(a) $\tilde{\Gamma}_2(2\mathbb{Z}[i])$ (b) $\tilde{\Gamma}_2(2\mathbb{Z}[i])$

Figure 12:

In order to solve the problem of describing a finite set of generators for a subgroup of finite index in $\mathcal{U}(\mathbb{Z}G)$ for an arbitrary nilpotent finite group G, it remains to deal with simple components $\mathbb{Q}Ge$ of $\mathbb{Q}G$ that are non-commutative division algebras D or $M(2, \mathcal{H}(a, b, \mathbb{Q}))$. As seen above, such a D is a classical quaternion algebra over $\mathbb{Q}(\xi_n)$ (with n a positive integer so that the order of 2 modulo n is odd) or over a real field $\mathcal{H}(\mathbb{Q}(\xi_n + \xi_n^{-1}))$. In the latter case, $\mathcal{H}(\mathbb{Q}(\xi_n + \xi_n^{-1}))$ is a totally definite quaternion algebra and hence (see [38, Lemma 21.3]) the unit group of an order $\emptyset$ in this algebra is central-by-finite. So, by the earlier remarks, if also $\mathbb{Q}Ge = D$, then the group $\langle b_{(n)} \mid b \text{ a Bass cylcic unit in } \mathbb{Z}G\rangle$ contains a subgroup of finite index in $\mathcal{U}(\emptyset)$.

Consequently, in case D is non-commutative division algebra that is a simple component of $\mathbb{Q}G$, for G nilpotent and finite, it remains to deal with orders $\mathcal{H}(\mathbb{Z}[\xi_n])$ in division algebras $\mathcal{H}(\mathbb{Q}(\xi_n))$. So far, this problem remains unsolved.

In [28], the author considers non-commutative division algebras D that split over $\mathbb{R}$. The non-split case is seemingly harder and even constructing sufficiently many explicit units is not trivial. For $\mathcal{H}(\mathbb{Z}[\sqrt{-d}])$ explicit constructions of units were given for all d in [24]. In Section 4 we also did this for other non-split division algebras.

A finite group G is said to be of Kleinian type if all Wedderburn components of $\mathbb{Q}G$ contain an order whose group of reduced norm one units acts discretely on $\mathbb{H}^3$. The classification of these groups was completed in [21]. An interesting consequence of this classification is that G is of Kleininan type if and only if $\mathcal{U}(\mathbb{Z}G)$ is virtually a direct product of free-by-free groups.

The following lemma follows, by the results of [18] mentioned before, if we also suppose that G is nilpotent.

Lemma 5.5 *Let G be a finite group and $A = \mathcal{H}(-1,\ -1,\ \mathbb{Q}[\sqrt{-d}])$, a non-split division algebra. Then A is not a Wedderburn component of $\mathbb{Q}G$.*

Proof. Our proof makes use of the classification of Amitsur [2] (see also [39, Theorems, 2.1.4, 2.1.5 and 2.1.6]) and results of Section 2.1 of [39].

Let $e \in \mathbb{Q}G$ be a primitive central idempotent and suppose that $\mathbb{Q}Ge = A$. Clearly A is not a totally definite quaternion algebra. We suppose first that Ge is not a Z-group. If $Ge = \mathrm{SL}(2, 5)$ or the binary octahedral group one would have that $\mathbb{Q}Ge$ is a totally definite quaternion algebra. If $g \in Ge$ is a central element of odd order m, say, then the center of A contains a root of unity and hence $d \in \{1, 3\}$, contradicting the fact that A is a division algebra. If $g \in Ge$ has odd order say m then $g + g^{-1}$ is central and so $\phi(m) = 4$, where ϕ is the Euler function. It follows that $m = 12$ and hence $d = 3$, a contradiction.

Suppose now that Ge is a Z-group. Let $1 \neq g \in Ge$ be of odd order, then g is non-central and $\mathbb{L} = \mathbb{Q}(\sqrt{-d}, g)$ is a maximal field. If $\mathbb{L} = \mathbb{Q}(g)$, then $\mathbb{Q}(g + g^{-1}) \cap \mathbb{R}$ is of index 2 in $\mathbb{L}$ and hence equals the center of A, a contradiction. Hence $[\mathbb{L} : \mathbb{Q}(\sqrt{-d})] = 2$, $\mathbb{Q}(g)$ is a quadratic extension of $\mathbb{Q}$ and thus $o(g) \in \{3, 4, 6\}$. It follows that $o(g) = 3$, $g + g^{-1}$ is central and hence $d = 3$, a contradiction. ∎

On the other hand $A = \mathcal{H}(-1,\ -1,\ \mathbb{Q}[\sqrt{-d}])$ is a Wedderburn component of $\mathbb{Q}(\sqrt{-d})Q_8 = 4\mathbb{Q}(\sqrt{-d}) \oplus \mathcal{H}(\mathbb{Q}(\sqrt{-d}))$. Hence, $\mathcal{U}(\mathbb{Z}[\sqrt{-d}]Q_8)$ contains $\mathcal{U}(\mathcal{H}(\mathbb{Z}[\sqrt{-d}])) \cap (1 + 2\mathcal{H}(\mathbb{Z}[\sqrt{-d}]))$ as a subgroup of finite index. We considered this group in case $d = 15$ but a set of generators is to big to be listed here.

Acknowledgment. The second author is grateful to the Vrije Universiteit Brussel for its hospitality during his sabbatical year, while this research was being done. The third author would like to thank Fred Simons for his assistance with Mathematica.

References

[1] I. Agol, M. Belolipetsky, P. Storm, K. Whyte, Finiteness of arithmetic hyperbolic reflection groups, Groups Geom. Dyn. 2 (2008), no. 4, 481-498.

[2] S. A. Amitsur, Finite subgroups of division rings, Trans. Amer. Math. Soc. 80 (1955), 361-386.

[3] A. Bak, U. Rehmann, The congruence subgroup and metacyclic problems for $SL_{n\geq 2}$ of division algebras, J. Algebra 78, 475-547 (1982).

[4] M.D. Baker, A.W. Reid, Arithmetic knots in closed 3-manifolds, Journal of Knot Theory and its Ramifications, Vol. 11, No. 6 (2002) 903-920.

[5] H. Bass, The Dirichlet Unit Theorem, Induced Characters and Whitehead Groups of Finite Groups, Topology 4 (1966), 391-410.

[6] H. Bass, J. Milnor, J.P. Serre, Solution of the congruence subgroup problem for SL_n $(n \geq 3)$ and Sp_{2n} $(n \geq 2)$, Publ. Math. I.H.E.S. **33**, 59-137 (1967).

[7] A. F. Beardon, The Geometry of Discrete Groups, Springer Verlag NY, 1983.

[8] M. R. Bridson, A. Haefliger, Metric Spaces of Non-Positive Curvature, Springer, Berlin, 1999.

[9] M.J. Jacobson, Jr., H.C. Williams, Solving the Pell Equation, CMS Books in Mathematics, Springer, New York, 2009.

[10] C. Corrales, E. Jespers, G. Leal, Á. del Río, Presentations of the Unit Group of an Order in a Non-Split Quaternion Algebra, Advances in Mathematics 186(2004), 498-524.

[11] A. Dooms, E. Jespers, A. Konovalov, From Farey symbols to generators for subgroups of finite index in integral group rings of finite groups, J. K-Theory 6 (2010), no. 2, 263-283.

[12] J, Elstrodt, F. Grunewald, J. Mennicke, Groups Acting on Hyperbolic Space, Springer Verlag, Berlin Heidelberg, 1998.

[13] E. Falbel, A Spherical CR structure on the compelement of the figure eight knot with discrete holonomy, J. Differential Geometry (79), 2008, 69-110.

[14] D. Flöge, Zur Struktur der PSL_2 über einigen imaginär-quadratischen Zahlringen. [On the structure of PSL_2 over some imaginary quadratic number rings.] Math. Z. 183 (1983), no. 2, 255-279.

[15] A. Giambruno, S.K. Sehgal, Generators of large subgroups of units of integral group rings of nilpotent groups, J. Algebra 174 (1995), 150-156.

[16] M. Gromov, Hyperbolic Groups, in Essays in Group Theory, M. S. R. I. Publ. 8, Springer, 1987, 75-263.

[17] E. Jespers, G. Leal, Generators of large subgroups of the unit group of integral group rings, Manuscripta Math. 78 (1993), no. 3, 303-315.

[18] E. Jespers, G. Leal, Degree 1 and 2 representations of nilpotent groups and applications to units of group rings, Manuscripta Mathematica 86 (1995), 479-498.

[19] E. Jespers, M. Parmenter, S.K. Sehgal, Central Units of Integral Group Rings of Nilpotent Groups, Proc. A.M.S. 124 (1996), no.4, 1007-1012.

[20] E. Jespers, G. Olteanu, A. del Rio, Rational group algebras of finite groups: from idempotents to units of integral group rings, Algebras and Representation Theory, to appear.

[21] E. Jespers, A. Pita, A. del Río, M. Ruiz, P. Zalesskii, Pavel Groups of units of integral group rings commensurable with direct products of free-by-free groups, Adv. Math. 212 (2007), no. 2, 692-722.

[22] S. Johansson, On fundamental domains of arithmetic Fuchsian groups, Math. Comp. 69 (2000),no.229, 339-349.

[23] S. O. Juriaans, I. B. S. Passi, D. Prasad, Hyperbolic Unit Groups, Proceedings of the American Mathematical Society, vol 133(2), 2005, $415 - 423$.

[24] S. O. Juriaans, I. B. S. Passi, A. C. Souza Filho, Hyperbolic Unit Groups and Quaternion Algebras, Proc. Indian Acad. Sci. (Math. Sci.), vol. 119, No. 1, February 2009, pp. 9-22.

[25] S. O. Juriaans, A. C. Souza Filho, Free Groups in Quaternion Algebras, Submitted.

[26] S. Katok, Reduction Theory for Fuchsian Groups, Math. Ann. 273, 461470 (1986).

[27] E. Kleinert, Units of classical orders: a survey, Enseign. Math. (2) 40 (1994), no. 3-4, 205-248.

[28] E. Kleinert, Units in skew fields, Progress in Math. 186, Birkhauser Verlag, Basel, 2000.

[29] G. S. Lakeland, Dirichlet-Ford Domains and Arithmetic Reflection Groups, Pacific J. Math., to appear.

[30] C. Maclachlan, A.W. Reid, The arithmetic of hyperbolic 3-manifolds, Graduate Texts in Mathematics, 219. Springer-Verlag, New York, 2003. xiv+463 pp.

[31] Makito Oichi, A fundamental polyhedron for the figure-eight knot group, Topology and its Applications 146-147 (2005) 15-19.

[32] D.S. Passman, Permutation Groups, Benjamin, New York (1968).

[33] J.G., Ratcliffe, Foundations of Hyperbolic Manifolds, Springer Verlag, New York, 1994.

[34] U. Rehmann, A survey of the congruence subgroup problem, in Algebraic K-theory, Lect. Notes Math. (966), 197-207 (1980).

[35] R. Riley, Applications of a Computer Implementation of Poincaré's Theorem on Fundamental Polyhedra, Mathematics of Computation, Vol. 40, No. 162 (Apr., 1983), pp. 607-632.

[36] R. Riley, A quadratic parabolic group, Math. Proc. Camb. Phil. Soc. (1973), 77, 281-288.

[37] R. Riley, An elliptical path from parabolic representations to hyperbolic structures, Topology of low-dimensional manifolds (Proc. Second Sussex Conf., Chelwood Gate, 1977), Lecture Notes in Math., 722, Springer, Berlin, 1979, pp. 99-133.

[38] S.K. Sehgal, Units in integral group rings, Longman Scientific & Technical, Pitman Monographs, Surveys in Pure and Applied Mathematics 69, 1993.

[39] M. Shirvani, B.A.F Wehrfritz, Skew linear groups, London Math. Soc., Lect. Notes Series 118, Cambridge Univ. Press, 1986.

[40] R. Swan, Generators and relations for certain special linear groups, Adv. in Math., v. 6, 1971, pp. 1-77.

[41] W.P. Thurston, Three-Dimensional Geometry and Topology, vol. 1, Princeton University Press, Princeton, NJ, 1997.

[42] L.N. Vaserstein, The structure of classic arithmetic groups of rank greater than one, Math. USSR Sbornik (20) 465-492 (1973).

[43] K. Vogtmann, Rational homology of Bianchi groups, Math. Annalen 272 (1985), 399-419.

Department of Mathematics,
Vrije Universiteit Brussel,
Pleinlaan 2, 1050 Brussel, Belgium
emails: efjesper@vub.ac.be and akiefer@vub.ac.be

Instituto de Matemática e Estatística,
Universidade de São Paulo (IME-USP),
Caixa Postal 66281, São Paulo,
CEP 05315-970 - Brasil
email: ostanley@usp.br

Departamento de Matematica
Universidade Federal da Paraiba
e-mail: andrade@mat.ufpb.br

Escola de Artes, Ciências e Humanidades,
Universidade de São Paulo (EACH-USP),
Rua Arlindo Béttio, 1000, Ermelindo Matarazzo, São Paulo,
CEP 03828-000 - Brasil
email: acsouzafilho@usp.br